

# A Novel Approach to Solve Gaussian Valued Neutrosophic Shortest Path Problems

Ranjan Kumar, S A Edalatpanah, Sripati Jha, Ramayan Singh

**Abstract:** We have exhibited a novel method for finding the neutrosophic shortest path problem (NSSPP) considering Gaussian valued Neutrosophic number. We have used linear programming approach for finding the NSSPP for Gaussian valued Neutrosophic number which is as per best of our information, hasn't been used till date for any other research work. In this article, we have introduced a novel method which deals Gaussian shaped neutrosophic problem easily without using any ranking method. We have used this method to solve uncertain network problems to find the shortest path which can help in taking crucial uncertain decisions. Finally, some numerical considerations are provided to show the effectiveness of the proposed model.

**Keyword:** Neutrosophic fuzzy numbers; shortest path problem; gaussian valued neutrosophic number; network; programming method.

## I. INTRODUCTION

A tool representing the partnership or relationship function is called a Fuzzy Set (FS) and handle the real world problems in which generally some type of uncertainty exists [1]. This concept was generalized by Atanassov [2] to intuitionistic fuzzy set (IFS) which is regarding membership function (MF) and non-membership function (NMF), the characteristic functions of the set. Moreover, Dutta and Ali [3] introduced gaussian fuzzy membership function. In 2017, Dutta and Limboo [4] extended this concept into soft set theory which is known as bell shaped fuzzy soft set. Beside this, several theories have been developed for uncertainties, including: Generalized orthopair FSs [5], Pythagorean FSs [6], picture FSs [7], Hesitant interval based neutrosophic linguistic sets [8], N-valued interval neutrosophic sets (NVINS) [9], Generalized Interval-Valued Triangular Intuitionistic FS [10], interval-valued Pythagorean FS [11], interval type 2 FSs [12]. In 1995, Smarandache [13] premised the theme of Neutrosophic sets (NS) and this is generalized from the FS [1] and IFS [2]. The NS are set of elements having a membership degree indeterminate-membership and also non membership with the criterion less than or equal to 3. Some extensions of NSs, including, bipolar NS [14], interval NS [15] single-valued NS [17], neutrosophic linguistic set [18], neutrosophic trapezoidal set [20], and triangular fuzzy neutrosophic set [21], are brought into picture to solve various problems.

**Manuscript published on 28 February 2019.**

\* Correspondence Author (s)

**Ranjan Kumar\***, Mathematics Department, National Institute of Technology, Adityapur, Jamshedpur, India.

**S A Edalatpanah**, Department of Industrial Engineering, Ayandegan Institute of Higher Education, Tonekabon, Iran

**Sripati Jha**, Mathematics Department, National Institute of Technology, Adityapur, Jamshedpur, India.

**Ramayan Singh**, Mathematics Department, National Institute of Technology, Adityapur, Jamshedpur, India.

© The Authors. Published by Blue Eyes Intelligence Engineering and Sciences Publication (BEIESP). This is an [open access](https://creativecommons.org/licenses/by-nc-nd/4.0/) article under the CC-BY-NC-ND license [http://creativecommons.org/licenses/by-nc-nd/4.0/](https://creativecommons.org/licenses/by-nc-nd/4.0/)

The neutrosophic number is an exceptional type of neutrosophic sets that extend the domain of numbers from those of real numbers to neutrosophic numbers. By generalizing SVNSSs [22], Wang et al. [22] premised the idea of IVNS. The IVNS [23] is a more general database to generalize the concept of different types of sets to express membership degrees truth, indeterminacy, and a false degree in terms of intervals. Thus several articles are published in the field of fuzzy and neutrosophic sets. However, to the best of our insight, very few strategies dealing exclusively with NSSPP.

Broumi et al. [24,25] first introduced a technique of finding SP under SV-trapezoidal and triangular fuzzy neutrosophic environment. Recently, Kumar et al. [26] suggested a new method which overcomes the shortcoming of Broumi et al. [24,25], proposed method. Broumi et al. [27] developed a new algorithm to solve SPP using bipolar neutrosophic setting. Additionally, Broumi et al. [28] also discussed an algorithmic approach based on a score function defined in [29] for solving NSPP on a network with IVNN as the edges but in some cases obtained data may not be single valued triangular or single valued trapezoidal neutrosophic numbers. For this purpose we introduced a novel method which deals with Gaussian shaped neutrosophic environment problem for finding SP and Gaussian valued neutrosophic shortest path length. Also we noticed that all of the above authors did not consider GVNSSPP to solve shortest path. As per all the available informations, there is no such method available for solving GVNSSPPs. This article is sectioned as: In Section 2, some basic information is provided along with detailed description of the concepts on neutrosophic set theory, and an arithmetic operation on the Gaussian valued neutrosophic numbers. Section 3, includes the existing linear programming method under crisp SPPs and it also includes a novel method for solving GVNSSPP. In Section 4, some numerical precedents are given to uncover the adequacy of the proposed model. Finally, the last section deals with the conclusion for all the proposed methodologies

## II. PRELIMINARIES

**Definition 2.1:** [22]: Let  $X$  be a space point or objects, with a genetic element in  $X$  denoted by  $x$ . A single-valued NS,  $V$  in  $X$  is characterised by three independent parts, namely truth-MF  $T_V$ , indeterminacy-MF  $I_V$  and falsity-MF  $F_V$ , such that  $T_V : X \rightarrow [0,1]$ ,  $I_V : X \rightarrow [0,1]$ , and  $F_V : X \rightarrow [0,1]$ .

Now,  $V$  is denoted as



$V = \{ \langle x, (T_V(x), I_V(x), F_V(x)) \rangle \mid x \in X \}$ , satisfying  
 $0 \leq T_V(x) + I_V(x) + F_V(x) \leq 3$ .

**Definition 2.2:** [30]:  $\hat{r}^N = \langle (\overline{\mu_{ij,l}}, \overline{\sigma_{ij,m}}), (\mu_{ij,l}, \sigma_{ij,m}), (\underline{\mu_{ij,l}}, \underline{\sigma_{ij,m}}) \rangle$  is a special NS on the real number set  $\mathbb{R}$ , whose truth-MF  $\Delta(x_i)$ , indeterminacy-MF  $\nabla(x_i)$ , and falsity-MF  $\nabla(x_f)$ , are given as follows:

$$\Delta(x_i) = \exp \left( -\frac{1}{2} \left( \frac{x - \overline{\mu_{ij,l}}}{\overline{\sigma_{ij,l}}} \right)^2 \right) \quad (1)$$

$$\nabla(x_i) = \exp \left( -\frac{1}{2} \left( \frac{x - \underline{\mu_{ij,l}}}{\underline{\sigma_{ij,l}}} \right)^2 \right) \quad (2)$$

$$\nabla(x_f) = \exp \left( -\frac{1}{2} \left( \frac{x - \underline{\mu_{ij,l}}}{\underline{\sigma_{ij,l}}} \right)^2 \right) \quad (3)$$

This definition is depicted in below figure I where truth function is shown by purple graph, falsity is shown by red graph and indeterminacy is shown by yellow color graph.




Figure 1. Graphical representation of GVNS number

**Definition 2.3:** [31,32]: Arithmetic operation). Let  $\hat{r}^N = \langle (\overline{r_{ij,l}}, \overline{r_{ij,m}}), (r_{ij,l}, r_{ij,m}), (\underline{r_{ij,l}}, \underline{r_{ij,m}}) \rangle$  and  $\hat{s}^N = \langle (\overline{s_{ij,l}}, \overline{s_{ij,m}}), (s_{ij,l}, s_{ij,m}), (\underline{s_{ij,l}}, \underline{s_{ij,m}}) \rangle$  be two arbitrary SVTNNs, and  $\theta \geq 0$ ; then:

$$\hat{r}^N \oplus \hat{s}^N = \langle (\overline{r_{ij,l} + s_{ij,l}}, \overline{r_{ij,m} + s_{ij,m}}), (r_{ij,l} + s_{ij,l}, r_{ij,m} + s_{ij,m}), (\underline{r_{ij,l} + s_{ij,l}}, \underline{r_{ij,m} + s_{ij,m}}) \rangle$$

$$\hat{r}^N \otimes \hat{s}^N = \langle (\overline{r_{ij,l} \cdot s_{ij,l}}, \overline{r_{ij,m} \cdot s_{ij,m}}), (r_{ij,l} \cdot s_{ij,l}, r_{ij,m} \cdot s_{ij,m}), (\underline{r_{ij,l} \cdot s_{ij,l}}, \underline{r_{ij,m} \cdot s_{ij,m}}) \rangle$$

if  $(\theta > 0)$

$$\theta \hat{r}^N = \langle (\theta \overline{r_{ij,l}}, \theta \overline{r_{ij,m}}), (\theta r_{ij,l}, \theta r_{ij,m}), (\theta \underline{r_{ij,l}}, \theta \underline{r_{ij,m}}) \rangle$$

**Definition 2.4:** [30] Let  $\hat{r}^N = \langle (\overline{r_{ij,l}}, \overline{r_{ij,m}}), (r_{ij,l}, r_{ij,m}), (\underline{r_{ij,l}}, \underline{r_{ij,m}}) \rangle$  then the score function is defined as follows:

$$s(\tilde{r}) = \frac{1}{6} \left[ 4 - (\overline{r_{ij,l}} - r_{ij,l} - \underline{r_{ij,l}}) + (\overline{r_{ij,m}} - r_{ij,m} - \underline{r_{ij,m}}) \right]$$

**Definition 2.5.** [30]: Let  $\hat{r}^N = \langle (\overline{r_{ij,l}}, \overline{r_{ij,m}}), (r_{ij,l}, r_{ij,m}), (\underline{r_{ij,l}}, \underline{r_{ij,m}}) \rangle$  and  $\hat{s}^N = \langle (\overline{s_{ij,l}}, \overline{s_{ij,m}}), (s_{ij,l}, s_{ij,m}), (\underline{s_{ij,l}}, \underline{s_{ij,m}}) \rangle$  be

two arbitrary SVTNNs, the ranking of  $\tilde{r}$  and  $\tilde{s}$  by score function is described as follows:

1. if  $s(\tilde{r}) > s(\tilde{s})$  then  $\tilde{r} \succ \tilde{s}$
2. if  $s(\tilde{r}) < s(\tilde{s})$  then  $\tilde{r} \prec \tilde{s}$

### III. THE PROPOSED METHOD

Before we start the main algorithm, we introduce existing crisp shortest path model.

#### A. Existing crisp model in SPP

In this section, we have a tendency to study the notation and existing linear model in crisp and proposed neutrosophic SPPs

#### Notations

$\Omega$  : Starting node

$\bar{\Omega}$  : Final destination node

$\sum_{k=1}^s x_{mk}$  : The total flow out of node  $s$ .

$\sum_{k=1}^s x_{mk}$  : The total flow into node  $s$ .

$RK_{mk}$  : The shortest distance from associate degree  $m^{\text{th}}$  node to  $k^{\text{th}}$  node.

The crisp SPP problem within the applied math model is as follows Kumar et al. [26]

$$\text{Min} = \sum_{m=1}^s \sum_{k=1}^s RK_{mk} \cdot x_{mk} \quad (4)$$

Subject to:

$$\sum_{m=1}^s x_{mk} - \sum_{k=1}^s x_{km} = \tilde{k}_m$$

for all  $x_{mk} \in \mathfrak{R}$  and non-negative where  $m, k = 1, 2, \dots, s$  and:



$$\tilde{\kappa}_m = \begin{cases} 1 & \text{if } m = \Omega, \\ 0 & \text{if } m = \Omega + 1, \Omega + 2, \dots, \Upsilon - 1 \\ -1 & \text{if } m = \Upsilon. \end{cases} \quad (5)$$

**B. Transformation of crisp SPP model into neutrosophic SPP**

If we tend to replace the parameter  $RK_{mk}$  into neutrosophic parameters, i.e..  $RK_{mk}^N$ , then the applied math model of the neutrosophic surroundings is as follows: Kumar et al. [26]

$$Min = \sum_{m=1}^s \sum_{k=1}^s RK_{mk}^N \cdot x_{mk} \quad (6)$$

Subject to:

$$\sum_{m=1}^s x_{mk} - \sum_{k=1}^s x_{km} = \tilde{\kappa}_m \quad m, k = 1, 2, \dots, s.$$

$x_{mk} \in \mathfrak{R}$  And are non-negative.

Step 2: use the arithmetic operation from the definition 2.3, then we get

$$GVC = \left\langle \left( \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq}, \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \right); \left( \sum_{p=1}^w \sum_{q=1}^w NGV_{pq,s} \cdot x_{pq}, \sum_{p=1}^w \sum_{q=1}^w NGV_{pq,t} \cdot x_{pq} \right); \left( \sum_{p=1}^w \sum_{q=1}^w \underline{NGV}_{pq,s} \cdot x_{pq}, \sum_{p=1}^w \sum_{q=1}^w \underline{NGV}_{pq,t} \cdot x_{pq} \right) \right\rangle \quad (9)$$

With subject to constraints (8)

Step 3: solve the following crisp SP problem using standard algorithm such as

$$\overline{NGV}_r^* = Min \overline{NGV}_r = \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,s} \cdot x_{pq} \quad (10)$$

With subject to constraints (8)

The optimum value of model (10), is  $\overline{NGV}_r^*$

Step 4: solve the following crisp SP problem using standard algorithm such as

$$\overline{NGV}_a^* = Min \overline{NGV}_a = \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \quad (11)$$

With subject to constraints

$$\sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq} = \overline{NGV}_r^*$$

Constraints of model (10)

The ideal value of equation (11) is  $\overline{NGV}_a^*$

Step 5: solve the following crisp SP problem using standard algorithm such as

$$NGV_r^* = Min NGV_r = \sum_{p=1}^w \sum_{q=1}^w NGV_{pq,s} \cdot x_{pq} \quad (12)$$

With subject to constraints

$$\sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} = \overline{NGV}_a^*$$

Constraints of model (11)

The ideal value of equation 12 is  $NGV_r^*$

**C. Algorithm: A novel approach for finding the SPP under GVNS environment**

In this section, we tend to provide a novel method for solving the NSP and the NSPL. For this purpose we consider GVNS numbers for the parameters.

Step 1: develop a GVNS model for solving SPP.

$$Min GVC = \left\langle \left( \overline{NGV}_{pq,s}, \overline{NGV}_{pq,t} \right), \left( NGV_{pq,s}, NGV_{pq,t} \right), \left( \underline{NGV}_{pq,s}, \underline{NGV}_{pq,t} \right) \right\rangle x_{pq} \quad (7)$$

Subject to:

$$\sum_{m=1}^s \overline{\pi}_{mk} - \sum_{k=1}^s \overline{\pi}_{mk} = \tilde{\kappa}_m$$

for all  $\overline{\pi}_{mk} \in \mathfrak{R}$  and non-negative where

$m, k = 1, 2, \dots, s$  and

$$\tilde{\kappa}_m = \begin{cases} 1 & \text{if } m = \Gamma, \\ 0 & \text{if } m = \Gamma + 1, \Gamma + 2, \dots, \Lambda - 1 \\ -1 & \text{if } m = \Lambda. \end{cases} \quad (8)$$

Step 6: Similarly, solve the following crisp SP problem using standard algorithm such as

$$NGV_a^* = Min NGV_a = \sum_{p=1}^w \sum_{q=1}^w NGV_{pq,t} \cdot x_{pq} \quad (13)$$

With subject to constraints

$$\sum_{p=1}^w \sum_{q=1}^w NGV_{pq,s} \cdot x_{pq} = NGV_r^*$$

Constraints of equation (12)

The ideal value of equation (13) is  $NGV_a^*$

Step 7: Again solve the following crisp SP problem using standard algorithm such as

$$\underline{NGV}_r^* = Min \underline{NGV}_r = \sum_{p=1}^w \sum_{q=1}^w \underline{NGV}_{pq,s} \cdot x_{pq} \quad (14)$$

With subject to constraints

$$\sum_{p=1}^w \sum_{q=1}^w NGV_{pq,t} \cdot x_{pq} = NGV_a^*$$

Constraints of equation (13)

The ideal value of equation 14 is  $\underline{NGV}_r^*$

Step 8: Again solve the following crisp SP problem using standard algorithm such as

$$\underline{NGV}_a^* = Min \underline{NGV}_a = \sum_{p=1}^w \sum_{q=1}^w \underline{NGV}_{pq,t} \cdot x_{pq} \quad (15)$$

With subject to constraints



$$\sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq} = \overline{NGV}_r^*$$

Constraints of equation (14)

The ideal value of equation (15) is  $\overline{NGV}_a^*$

**Theorem 1:** The ideal value of equation (15) provides the optimum value of IVNSSP problem (7).

**Proof:** let  $x_{pq}^*$  be the ideal arrangement of equation (18) and  $\tilde{x}_{pq}$  be an arbitrary neutrosophic viable solution of IVNSSP (7). the solution system of the proposed technique confirms that the most excellent solution of problem (15) is the greatest answer of the problems (7)-(14). Owing the optimality of  $x_{pq}^*$  for problem (10) and feasibility of  $\tilde{x}_{pq}$  for problem (10), we conclude that

$$\overline{NGV}_r^* = \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,s} \cdot x_{pq}^* \leq \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,s} \cdot \tilde{x}_{pq}$$

owing the optimality of  $x_{pq}^*$  for hassle (11) and we conclude that

$$\overline{NGV}_a^* = \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,t} \cdot x_{pq}^* \leq \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,t} \cdot \tilde{x}_{pq} \quad \text{similar}$$

discussions make sure that

$$\overline{NGV}_r^* = \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq} \leq \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,s} \cdot \tilde{x}_{pq} ;$$

$$\overline{NGV}_a^* = \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \leq \sum_{r=1}^w \sum_{m=1}^w \overline{NGV}_{pq,t} \cdot \tilde{x}_{pq} ;$$

$$\overline{NGV}_r^* = \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq} \leq \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot \tilde{x}_{pq} ;$$

$$\overline{NGV}_a^* = \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \leq \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot \tilde{x}_{pq} ;$$

Therefore

$$GVC = \left\langle \begin{matrix} \left( \overline{NGV}_r^*, \overline{NGV}_a^* \right) \\ \left( \overline{NGV}_r^*, \overline{NGV}_a^* \right) \\ \left( \overline{NGV}_r^*, \overline{NGV}_a^* \right) \end{matrix} \right\rangle \leq \left\langle \begin{matrix} \left( \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq}, \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \right) \\ \left( \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq}, \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \right) \\ \left( \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,s} \cdot x_{pq}, \sum_{p=1}^w \sum_{q=1}^w \overline{NGV}_{pq,t} \cdot x_{pq} \right) \end{matrix} \right\rangle$$

#### IV. EXAMPLE OF NETWORK

To justify our proposed algorithms, we consider a network shown in fig. 3 Kumar et al. [33]




Figure 2. A network with eleven vertices and twenty-five edges Kumar et al. [33]

**Example 4.1.** Consider **Figure 2** with 11 nodes and 25 edges, where SV is 1 and DV is 11. The GVNS time is given in Table 1.

**I. Table 2. The conditions of Example 4.1.**

T	H	TrNS time	T	H	TrNS time
1	2	<(15,3);(17,5);(8,1)>	9	7	<(25,5);(20,6);(9,7)>
1	6	<(14,2);(5,1);(11,6)>	1	7	<(28,8);(6,1);(16,2)>
1	9	<(15,3);(4,1);(12,5)>	4	6	<(19,10);(8,3);(10,5)>
1	1	<(4,1); (5,1);(6,2)>	1	1	<(50,11);(10,6);(12,0)>
2	3	<(6,4);(5,3);(3,1)>	1	3	<(3,1);(2,1);(2,1)>
2	5	<(16,8);(11,7);(7,2)>	4	1	<(31,20);(9,7);(17,1)>
3	4	<(24,2);(8,1);(15,9)>	2	9	<(30,5);(15,5);(16,6)>
3	5	<(27,9);(9,6);(15,9)>	3	8	<(38,30);(20,10);(27,20)>
4	5	<(25,4);(13,2);(12,8)>	6	1	<(35,21);(9,7);(18,1)>
5	6	<(24,6);(13,2);(13,2)>	7	1	<(13,3);(3,1);(2,1)>
7	6	<(23,3);(11,4);(15,7)>	9	8	<(19,9);(8,7);(7,4)>
8	4	<(28,9);(15,7);(16,10)>	9	1	<(26,4);(5,3)(18,8)>
8	7	<(30,20);(8,6);(19,9)>		0	

**Solution:** Applying steps 1-8 in proposed Algorithm, the GVNSSP is 1 → 10 → 7 → 11 and the GVNSSPL is < (45, 12); (14, 3); (24, 5

>. The result is shown in below figure 3 and we can conclude that the GVNSSPL preserve the GVNS structure. Moreover, the suggested shortest route is depicted in figure 4.






Figure 3. the optimum solution also preserve the weight of gaussian fuzzy structure.

Proposed algorithm is executed numerically as follows:

Step 1: The GVNSSP model is detailed as shown in equation 7:

$$\begin{aligned} \text{Min GVC} = & \langle (15,3);(17,5);(8,1) \rangle \cdot x_{12} + \langle (14,2);(5,1);(11,6) \rangle \cdot x_{16} + \langle (3,1);(2,1);(2,1) \rangle \cdot x_{13} \\ & + \langle (15,3);(4,1);(12,5) \rangle \cdot x_{19} + \langle (4,1);(5,1);(6,2) \rangle \cdot x_{110} + \langle (6,4);(5,3);(3,1) \rangle \cdot x_{23} + \\ & \langle (25,5);(20,6);(9,7) \rangle \cdot x_{97} + \langle (28,8);(6,1);(16,9) \rangle \cdot x_{107} + \langle (19,10);(8,3);(10,5) \rangle \cdot x_{46} + \\ & \langle (50,11);(10,6);(12,7) \rangle \cdot x_{1011} + \langle (16,8);(11,7);(7,2) \rangle \cdot x_{25} + \langle (31,20);(9,7);(17,14) \rangle \cdot x_{411} + \\ & \langle (24,2);(8,1);(15,9) \rangle \cdot x_{34} + \langle (30,5);(15,5);(16,6) \rangle \cdot x_{29} + \langle (27,9);(9,6);(15,9) \rangle \cdot x_{35} + \\ & \langle (38,30);(20,10);(27,20) \rangle \cdot x_{35} + \langle (25,4);(13,2);(12,8) \rangle \cdot x_{45} + \langle (35,21);(9,7);(18,10) \rangle \cdot x_{611} + \\ & \langle (24,6);(13,2);(13,2) \rangle \cdot x_{56} + \langle (13,3);(3,1);(2,1) \rangle \cdot x_{711} + \langle (23,3);(11,4);(15,7) \rangle \cdot x_{76} + \\ & \langle (19,9);(8,7);(7,4) \rangle \cdot x_{98} + \langle (28,9);(15,7);(16,10) \rangle \cdot x_{84} + \langle (26,4);(5,3);(18,8) \rangle \cdot x_{910} + \\ & \langle (30,20);(8,6);(19,9) \rangle \cdot x_{87} \end{aligned}$$

Subject to constraints in line with equation (8)

$$\begin{aligned} x_{12} + x_{13} + x_{16} + x_{19} + x_{110} &= 1; & x_{23} + x_{25} + x_{29} &= x_{12}; & x_{34} + x_{35} + x_{38} &= x_{13} + x_{23} \\ x_{45} + x_{46} + x_{411} &= x_{34} + x_{84} & x_{56} &= x_{25} + x_{35} + x_{45}; & x_{611} &= x_{56} + x_{46} + x_{16} + x_{76}; \\ x_{1011} &= x_{110} + x_{910}; & x_{76} + x_{711} &= x_{87} + x_{97} + x_{107}; & x_{84} + x_{87} &= x_{38} + x_{98}; \\ x_{98} + x_{97} + x_{910} &= x_{19} + x_{29}; & x_{1011} &= x_{110} + x_{910}; & x_{411} + x_{611} + x_{711} + x_{1011} &= 1; \end{aligned}$$

Step 2: Execute the arithmetic operation on equation (7) then proceed to step 3

Step 3: Now we get the linear standard equation (10)

$$\begin{aligned} \overline{NGV}_r^* = \text{Min } \overline{NGV}_r = & 15 \cdot x_{12} + 14 \cdot x_{16} + 15 \cdot x_{19} + 15 \cdot x_{110} + 6 \cdot x_{23} + 16 \cdot x_{25} + 24 \cdot x_{34} + 27 \cdot x_{35} + 25 \cdot x_{45} + \\ & 24 \cdot x_{56} + 23 \cdot x_{76} + 28 \cdot x_{84} + 30 \cdot x_{87} + 25 \cdot x_{97} + 28 \cdot x_{107} + 19 \cdot x_{46} + 50 \cdot x_{1011} + 3 \cdot x_{13} + \\ & 31 \cdot x_{411} + 30 \cdot x_{29} + 38 \cdot x_{38} + 35 \cdot x_{611} + 13 \cdot x_{711} + 19 \cdot x_{98} + 26 \cdot x_{910} \end{aligned}$$

Subject to constraints in line with equation (8)

After executing the LPP then the optimal solution is 45.

Step 4: Now we get the linear standard equation (11)



$$\overline{NGV}_a^* = \text{Min } \overline{NGV}_a = 3 \cdot x_{12} + 2 \cdot x_{16} + 3 \cdot x_{19} + 1 \cdot x_{110} + 4 \cdot x_{23} + 8 \cdot x_{25} + 2 \cdot x_{34} + 9 \cdot x_{35} + 4 \cdot x_{45} + 6 \cdot x_{56} + 3 \cdot x_{76} + 20 \cdot x_{84} + 9 \cdot x_{87} + 5 \cdot x_{97} + 8 \cdot x_{107} + 10 \cdot x_{46} + 11 \cdot x_{1011} + 1 \cdot x_{13} + 30 \cdot x_{411} + 5 \cdot x_{29} + 20 \cdot x_{38} + 21 \cdot x_{611} + 3 \cdot x_{711} + 9 \cdot x_{98} + 4 \cdot x_{910}$$

Subject to constraints in line with equation (10) and

$$15 \cdot x_{12} + 14 \cdot x_{16} + 15 \cdot x_{19} + 15 \cdot x_{110} + 6 \cdot x_{23} + 16 \cdot x_{25} + 24 \cdot x_{34} + 27 \cdot x_{35} + 25 \cdot x_{45} + 24 \cdot x_{56} + 23 \cdot x_{76} + 28 \cdot x_{84} + 30 \cdot x_{87} + 25 \cdot x_{97} + 28 \cdot x_{107} + 19 \cdot x_{46} + 50 \cdot x_{1011} + 3 \cdot x_{13} + 31 \cdot x_{411} + 30 \cdot x_{29} + 38 \cdot x_{38} + 35 \cdot x_{611} + 13 \cdot x_{711} + 19 \cdot x_{98} + 26 \cdot x_{910} = 45;$$

After execute the LPP then the optimal solution is 12

Similarly proceed from step 5 to step 8, we get the final optimum solution the NSSP is 1 → 10 → 7 → 11 and the NSSPL is <(45,12);(14,3);(24,5)>. Finally the shortest route is shown in figure 4:




Figure 4: shown the suggested shortest route

V. CONCLUSION

Conventional SP issue expects exact qualities for the curve loads which aren't generally the situation in genuine circumstances. In this article, briefest issue for numbering esteemed neutrosophic circular segment loads has been explored. We initially figured the issue in the number esteemed neutrosophic condition. At that point, we proposed another arrangement approach for understanding whole number esteemed neutrosophic most limited issue. We have changed over the GVNSSPP issue under thought to multi-objective LP issues which can be illuminated utilizing the standard LP calculations. According to the proposed optimization manner, the Gaussian-valued neutrosophic source weight has preserved the shape of a Gaussian-valued neutrosophic quantity. The numerical results show that the new algorithms outperform the present day strategies. In future, we will extend the method to more complicated community issues involving Gaussian-valued neutrosophic costs.

REFERENCES

1. R. Kumar, S. Jha, and R. Singh, "A different approach for solving the shortest path problem under mixed fuzzy environment," *International journal of fuzzy system applications*, vol. 9, issue 2 p. Article 6, Forthcoming.
2. M. Abdel-Basset, M. Gunasekaran, M. Mohamed, and F. Smarandache, "A novel method for solving the fully neutrosophic linear programming problems," *Neural Computing and Applications*, Mar. 2018.

3. M. Mohamed, M. Abdel-Basset, A. N. Zaid, and F. Smarandache, "Neutrosophic integer programming problem," *Neutrosophic Sets and Systems*, vol. 15, pp. 3-7, 2017.
4. F. Karaaslan, "Gaussian single-valued neutrosophic numbers and its application in multi-attribute decision making," *Neutrosophic Sets and Systems*, vol. 22, pp. 101-117, 2018.
5. R. Şahin and P. Liu, "Maximizing deviation method for neutrosophic multiple attribute decision making with incomplete weight information," *Neural Computing and Applications*, vol. 27, pp. 2017-2029, Oct. 2016.
6. S. Broumi, A. Bakali, M. Talea, F. Smarandache, and M. Ali, "Shortest path problem under interval valued neutrosophic setting," *Journal of Fundamental and Applied Sciences*, vol. 10, pp. 168-174, 2018.
7. S. Broumi, A. Bakali, M. Talea, F. Smarandache, and M. Ali, "Shortest path problem under bipolar neutrosophic setting," in *Advanced Research in Area of Materials, Aerospace, Robotics and Modern Manufacturing Systems*, vol. 859, Feb. 2017, pp. 59-66.
8. R. Kumar, S. A. Edalatpanah, S. Jha, S. Broumi, and A. Dey, "Neutrosophic shortest path problem," *Neutrosophic Sets and Systems*, vol. 23, pp. 5-15, 2018.
9. S. Broumi, A. Bakali, M. Talea, F. Smarandache, and L. Vladareanu, "Computation of shortest path problem in a network with SV-trapezoidal neutrosophic numbers," in *2016 International Conference on Advanced Mechatronic Systems (ICAMechS)*, 2016, pp. 417-422.
10. S. Broumi, A. Bakali, M. Talea, F. Smarandache, and L. Vladareanu, "Computation of Shortest Path Problem in a Network with SV-Triangular Neutrosophic Numbers," in *IEEE International Conference on Innovations in Intelligent Systems and Applications (INISTA)*, Gdynia, Poland, 2017, pp. 426-431.
11. H. Wang, F. Smarandache, Y-Q Zhang, and R. Sunderraman, *Interval Neutrosophic Sets and Logic: Theory and Applications in Computing*. Infinite Study, 2005.
12. H. Wang, F. Smarandache, Y-Q Zhang, and R. Sunderraman, "Single valued neutrosophic sets," *Multispace and Multistructure*, 2005.
13. P. Biswas, S. Pramanik, and B. C. Giri, "Aggregation of triangular fuzzy neutrosophic set information and its application to multi-attribute decision making," *Neutrosophic Sets and Systems*, vol. 12, pp. 20-40, 2016.
14. Pramanik, and B. C. Giri, "TOPSIS Strategy for Multi-Attribute Decision Making with Trapezoidal Neutrosophic Numbers," *Neutrosophic Sets and Systems*, vol. 19, pp. 29-39, 2018.
15. P. Biswas, S.
16. J. Ye, "An extended TOPSIS method for multiple attribute group decision making based on single valued neutrosophic linguistic numbers," *Journal of Intelligent & Fuzzy Systems*, vol. 28, pp. 247-255, 2015.
17. P. Liu, "The aggregation operators based on archimedean t-conorm and t-norm for single-valued neutrosophic numbers and their application to decision making," *International Journal of Fuzzy Systems*, vol. 18, pp. 849-863, 2016.
18. J. Ye, "Similarity measures between interval neutrosophic sets and their applications in multicriteria decision-making," *Journal of Intelligent & Fuzzy Systems*, vol. 26, pp. 165-172, 2014.



19. Uluçay, I. Deli, and M. Şahin, "Similarity measures of bipolar neutrosophic sets and their application to multiple criteria decision making," *Neural Computing and Applications*, vol. 29, pp. 739-748, 2018.
20. F. Smarandache, , Minh Perez, Ed.: American Research Press, 1999.
21. J. M. Mendel, R. I. John, and F. Liu, "Interval type-2 fuzzy logic systems made simple," *IEEE Transactions on Fuzzy Systems*, vol. 14, pp. 808-821, 2006.
22. H. Garg, "A novel accuracy function under interval-valued Pythagorean fuzzy environment for solving multicriteria decision making problem," *Journal of Intelligent & Fuzzy Systems*, vol. 31, pp. 529-540, 2016.
23. P. Dutta and P. Talukdar, "A novel arithmetic technique for generalized interval-valued triangular intuitionistic fuzzy numbers and its application in decision making," *The Open Cybernetics & Systemics Journal*, vol. 12, pp. 72-120, 2018.
24. S. Broumi, I. Deli, and F. Smarandache, "N-Valued Interval Neutrosophic Sets and Their Application in Medical Diagnosis," *Critical Review : Center for Mathematics of Uncertainty, Creighton University , Omaha, NE, USA.*, vol. 10, pp. 45-69, 2015.
25. J. Ye, "Hesitant interval neutrosophic linguistic set and its application in multiple attribute decision making," *International Journal of Machine Learning and Cybernetics*, pp. 1-12, Nov. 2017.
26. C. Cùròng, "Picture fuzzy sets," *Journal of Computer Science and Cybernetics*, vol. 30, pp. 409-420, 2014.
27. R. R. Yager, "Pythagorean fuzzy subsets," in *2013 Joint IFSA World Congress and NAFIPS Annual Meeting (IFSA/NAFIPS)*, Edmonton, AB, Canada, 2013, pp. 57-61.
28. R. R. Yager, "Generalized Orthopair Fuzzy Sets," *IEEE Transactions on Fuzzy Systems*, vol. 25, pp. 1222-1230, 2017.
29. P. Dutta and B. Limboo, "Bell-shaped fuzzy soft sets and their application in medical diagnosis," *Fuzzy Information and Engineering*, vol. 9, pp. 67-91, 2017.
30. P. Dutta and T. Ali, "Uncertainty modelling in risk analysis: A fuzzy set approach," *International Journal of Computer Applications*, vol. 43, pp. 35-39, 2012.
31. K. Atanassov, "Intuitionistic fuzzy sets," *Fuzzy Sets and Systems*, vol. 20, p. 87, 1986.
32. L. A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, pp. 338-358, 1965.

#### AUTHORS PROFILE:



**Ranjan Kumar**, Research Scholar, National Institute of Technology, Jamshedpur. His fields of interest are uncertainty theory, Fuzzy Optimization and Operational Research. He is a Member of renowned societies such as IAENG Society of Operations Research and European Operational Research Societies (EURO ).



**Dr. S A Edalatpanah**. Received his PhD in applied mathematics from the University of Guilan, Rasht, Iran. He is an academic member of Guilan University, Islamic Azad University, and Ayandegan Institute of Higher Education. His fields of interest are numerical linear algebra, soft computing and optimization. He is Editor-in-Chief of the International Journal of Research in Industrial Engineering at [www.riejournal.com](http://www.riejournal.com).



**Dr. Sripati Jha**, Associate Proffesor, National Institute of Technology, Jamshedpur. His fields of interest are Fuzzy , Fuzzy Optimization and Operational Research. He is serving as Dean Alumni & Industrial Relations which is highly respectable administrative position in the institutaion. He has also served as Dean Student Welfare in the past. He has succesfully guided one research scholar to complete the doctorate degree and three scholars are undergoing their doctarate degree under his supervision. Additionally, he has also provided guidance for projects related to Master degree in Mathematics.



**Dr. Ramayan Singh**, Associate Proffesor, National Institute of Technology, Jamshedpur. His fields of interest are Uncertainty theory, and fluid dynamics and numerical and computation mathematics.