

A Research of Achromatic and B-Chromatic Number of Multistar-Related Graphs

K.P.Thilagavathy, A.Santha, K.Paramasivam

ABSTRACT--- *The structural properties, achromatic number and b-chromatic number of central graph of double star graph and triple star graph have been studied in [6]. In this paper, these studies have been carried out, for the central graph of any multi star graph and central graph of shadow graph of star graph and double star graph.*

Key Words - Double Star graph, Triple Star graph, multi star graph, achromatic number, b-chromatic number, Shadow graph
Mathematics subject classification: 05C15

I. INTRODUCTION

Graph theory and graph colouring have wide applications in physics, chemistry, social sciences, networking, VLSI circuits [4] and many other fields.

A Star graph a_n is the bi partite graph $K_{1,n}$. The multistar graph $K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m})$ is the graph got by adding n_1, n_2, \dots, n_m leaves respectively to the m vertices of the complete graph K_m . The number of spanning trees of a specific family of multistar graph was studied by Nikolopoulos and Rondogiannis [3].

Consider a simple undirected graph G . To form its central graph $C(G)$ we introduce a new node on every edge of G and join the non-adjacent nodes of G . In 1999, Irving and Manlove introduced the idea of b-chromatic number, which is the maximum value of k for which G has a proper colouring with k colours in such a way that every colour is adjacent to every other colour.

The achromatic number was introduced by Harary[1]. An achromatic colouring is a proper vertex colouring such that each pair of colours is adjacent by at least one edge. The largest possible number of colours in an achromatic colouring of G is called the achromatic number of G and it is denoted by $\psi(G)$.

The concept of generalisation of achromatic colouring of graphs was studied by Roopesh and Thilagavathi[5], who proved that the achromatic number of the central graph of any simple graph $G(V,E)$ with $|V| = n$ is less than or equal to $n + 1$. The achromatic and b-chromatic number of various graphs have been studied in [6,7,8]

The Shadow graph $D_2(G)$, of a connected graph G is constructed by taking two copies of G , say G' and G'' . Join

each vertex u' in G' to the neighbours of the corresponding vertex v'' in G'' . The following results have been discussed in [8].

The Structural Properties and the colouring of Central graph of Double Star Graph

1. The number of vertices in the graph $C[K_2(a_n, a_r)]$ is $p = 2(n + r) + 3$
2. The number of edges in the graph $C[K_2(a_n, a_r)]$ is $q = \frac{(n + r + 1)(n + r + 4)}{2}$
3. The maximum degree in the graph $C[K_2(a_n, a_r)]$ is $\Delta = n + r + 1$
4. For any Double Star graph $K_2(a_n, a_r)$ the achromatic number $\psi[C(K_2(a_n, a_r))] = n + r + 2$
5. For any Double Star graph $K_2(a_n, a_r)$ the b-chromatic number $\varphi[C(K_2(a_n, a_r))] = n + r + 1$

The Structural Properties and the colouring of Central graph of Triple Star Graph

1. The number of vertices in the graph $C[K_3(a_n, a_r, a_t)]$ is $p = 2(n + r + t + 3)$
2. The number of edges in the graph $C[K_3(a_n, a_r, a_t)]$ is $q = \frac{(n+r+t+3)(n+r+t+4)}{2}$
3. The maximum degree in the graph $C[K_3(a_n, a_r, a_t)]$ is $\Delta = n + r + t + 2$
4. For any Triple Star graph $K_3(a_n, a_r, a_t)$ the b-chromatic number $\varphi[C(K_3(a_n, a_r, a_t))] = n + r + t + 1$
5. For any Triple Star graph $K_3(a_n, a_r, a_t)$ the achromatic number $\psi[C(K_3(a_n, a_r, a_t))] = n + r + t + 3$

The generalisation of the above results to multi star graphs is discussed here.

The Achromatic and b-chromatic number of Central graph of Multi Star graph

Theorem 2.1

The achromatic number of central graph of multi star graph is $\psi[C(K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m}))] = p$, where p denotes the number of vertices in the multi star graph.

Proof:

Let $K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m})$ be a multi star graph formed by adding n_1, n_2, \dots, n_m leaves to the m nodes of K_m . Let p be the total number of vertices in the multi star graph. That is $p = n_1 + n_2 + \dots + n_m + m$. In


Revised Manuscript Received on August 14, 2019.

K.P.Thilagavathy, Assistant Professor, Department of Science and Humanities, Kumaraguru College of Technology, Coimbatore-641049, T.N. India

A.Santha, Associate Professor, Department of Science and Humanities, Kumaraguru College of Technology, Coimbatore-641049, T.N. India

K.Paramasivam, Professor, Department of Electrical and Electronics Engineering, Kumaraguru College of Technology, Coimbatore-641049, T.N. India

$C(K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m}))$, the $p - m$ leaf vertices are given $p - m$ different colours and the remaining m vertices of the complete graph are coloured by m other colours. Hence $(p - m) + m$ colours are needed to colour this graph and by this construction it is the maximal one, and it is achromatic.



Example-1

$$\psi [C(K_4(a_{3_1}, a_{3_2}, a_{3_3}, a_{2_4}))] = 15$$

RESULTS

Theorem 2.2


The b-chromatic number of central graph of multi star graph

$$\varphi [C(K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m}))] = \begin{cases} p - \binom{m}{2}, & m = \text{even} \\ p - \binom{m+1}{2}, & m = \text{odd} \end{cases}$$

where p denotes the number of vertices in the multi star graph.

Proof:

Let $K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m})$ be a multi star graph formed by adding n_1, n_2, \dots, n_m leaves to the m nodes of K_m . Let p be the total number of vertices in the multi star graph. That is $p = n_1 + n_2 + \dots + n_m + m$. In $C[K_m(a_{n_1}, a_{n_2}, \dots, a_{n_m})]$ we need $p - m$ colours to colour the leaf vertices of $a_{n_1}, a_{n_2}, \dots, a_{n_m}$. Then the complete graph K_m can be coloured by $\frac{m}{2}$ colours if m is even. The same colour can be given to two adjacent vertices. In the case when m is odd we use $\frac{m+1}{2}$ colours to colour $m + 1$ vertices of K_m as in the above method. The last vertex can be coloured using any of the $\frac{m+1}{2}$ colours. This colouring is achromatic and it is the maximal one.



Example-2 Example-3

$$\varphi [C(K_5(a_{1_1}, a_{2_1}, a_{3_2}, a_{4_2}, a_{5_2}))] = 9$$

$$\varphi [C(K_4(a_{1_2}, a_{2_2}, a_{3_2}, a_{4_2}))] = 10$$

3. The Achromatic and b-chromatic number of Central graph of Shadow graph of Star graphs

The Structural properties of central graph of Shadow graph of Star graphs

1. The number of vertices in the graph $C(D_2(K_{1,n}))$ is $p = 6n + 2$
2. The number of edges in the graph $C(D_2(K_{1,n}))$ is $q = 2n^2 + 7n + 1$
3. The maximum degree in the graph $C(D_2(K_{1,n}))$ is $\Delta = 2n + 1$


Theorem 3.1

The achromatic number of central graph of $D_2(K_{1,n})$ is $\psi [C(D_2(K_{1,n}))] = 2(n + 1)$

Proof: Let $K_{1,n}$ be the star graph. In $D_2(K_{1,n})$, let v_0, v_1, \dots, v_n be the vertices of the first star graph and u_0, u_1, \dots, u_n be the corresponding vertices of the second star graph. By the definition of the shadow graph all the v_i 's are connected with u_0 and all the u_i 's are connected with v_0 . In $C(D_2(K_{1,n}))$, subdivide all the edges of $D_2(K_{1,n})$ once and join all the non-adjacent vertices.

Let u_0v_i be the newly introduced vertex on the edge joining the vertices u_0 and v_i and v_0u_i be the vertex on the edge joining the vertices v_0 and u_i where $i = 1, 2, 3 \dots n$. Let v_0v_i and u_0u_i be the newly introduced vertices on the edges joining the vertices v_0, v_i and u_0, u_i respectively where $i = 1, 2, 3 \dots n$. Consider the two sets of colours $C = \{C_0, C_1, \dots, C_n\}$ and $C' = \{C'_0, C'_1, \dots, C'_n\}$. Assign C_i to v_i and C'_i to u_i where $i = 0, 1, 2, 3 \dots n$.

Hence the pairs of colours (C_i, C_j) and (C'_i, C'_j) where $i \neq j, j \neq 0$ are adjacent. The pair (C_0, C'_0) is adjacent. For making the remaining pairs of colours adjacent for an achromatic colouring, consider the following procedure: For $1 \leq i \leq n$ assign C_0 to the vertices u_0v_i and u_0u_i . For $1 \leq i \leq n$ assign C'_0 to the vertices v_0v_i and v_0u_i . By this construction any pair of colours in $C \cup C'$ is adjacent by at least one edge and this colouring is the maximal possible one. Hence $\psi [C(D_2(K_{1,n}))] = 2(n + 1)$.



Example-4 $\psi [C(D_2(K_{1,3}))] = 8$

Theorem 3.2

The b-chromatic number of central graph of $D_2(K_{1,n})$ is $\varphi [C(D_2(K_{1,n}))] = 2n + 1$.

Proof: Let $K_{1,n}$ be the star graph. In $D_2(K_{1,n})$, let v_0, v_1, \dots, v_n be the vertices of the first star graph and u_0, u_1, \dots, u_n be the corresponding vertices of the second star graph. By the definition of the shadow graph all the v_i 's are connected with u_0 and all the u_i 's are connected with v_0 . In $C(D_2(K_{1,n}))$, subdivide all the edges of $D_2(K_{1,n})$ once and join all the non-adjacent vertices.

Let u_0v_i be the newly introduced vertex on the edge joining the vertices u_0 and v_i and v_0u_i be the vertex on the edge joining the vertices v_0 and u_i where $i = 1, 2, 3 \dots n$. Let v_0v_i and u_0u_i be the newly introduced vertices on the edges joining the vertices v_0, v_i and u_0, u_i respectively where $i = 0, 1, 2, 3 \dots n$.


Consider $v_i, i = 1, 2, 3, \dots, n$. In $C(D_2(K_{1,n}))$, v_i is adjacent to all other v_j 's, ($j \neq 0$) and all u_i 's. Similarly $u_i, i = 1, 2, 3, \dots, n$ is adjacent to all other u_j 's, ($j \neq 0$) and v_i 's, $i = 1, 2, 3 \dots, n$. Also v_0 is adjacent only to u_0 and u_0 is adjacent only to v_0 . Consider the two sets of colours $C = \{C_0, C_1, \dots, C_n\}$ and $C' = \{C'_1, \dots, C'_n\}$. To make the colouring as b-chromatic consider the following colouring procedure:

- For $1 \leq i \leq n$ assign C'_i to the vertices v_0v_i
- For $1 \leq i \leq n$ assign C_i to the vertices v_0u_i

- For $1 \leq i \leq n$ assign C_0 to the vertices u_0v_i and assign C_0 to the vertex u_0u_1

Here the representative vertices of the $2n + 1$ colour classes that are adjacent to a vertex in every other class are $v_0, v_1, \dots, v_n, u_1, \dots, u_n$.

Hence $\varphi [C(D_2(K_{1,n}))] = 2n + 1$.



Example-5 $\varphi [C(D_2(K_{1,3}))] = 7$

4. The Achromatic and b-chromatic number of Central graph of Shadow graph of Double Star graph

The Structural properties of central graph of Shadow graph of Double Star graph

1. The number of vertices in $D_2(K_2(a_n, a_r))$ is $p = 2(2 + n + r)$
2. The number of edges in $D_2(K_2(a_n, a_r))$ is $q = 4(n + r) + 2$

Observation 4.1

1. The achromatic number of central graph of $D_2(K_2(a_n, a_r))$ is $\psi [C(D_2(K_2(a_n, a_r)))] = 2(n + r + 2)$
2. The b-chromatic number of central graph of $D_2(K_2(a_n, a_r))$ is $\varphi [C(D_2(K_2(a_n, a_r)))] = 2(n + r)$

II. CONCLUSION

In this research work, the achromatic and b-chromatic numbers of central graph of double star, triple star and any multi star graph have been discussed. Also, the structural properties, the achromatic and b-chromatic numbers of the central graphs of shadow graph of star graph and double star graph have been studied.

REFERENCES

1. Harary, F. Graph Theory, Narosa Publishing 1969.
2. Irving, R.W and Manlove, D.F. "The b-Chromatic number of a graph", Discrete Applied Mathematics 91(1-3) 127-141, 1999.
3. Nikipoulos, S.D and Rondogiannis, P. "On the Number of Spanning trees of Multistar Related Graphs", Volume 65, Issue 4, 183-188, 1998.
4. Paramasivam, K, Gunavathi, K, Sathishkumar, P, "Graph Theory Based Approach for Low Power Combinational Circuit Testing", proceedings of International AMSE conference MS2004, Lyon, France, July 5-6, 2004.



5. Thilagavathi, K, Roopesh, N. '*Generalisation of achromatic colouring of central graphs*', Electronic Notes in Discrete Mathematics, vol. 33, pp. 147-152, 2009.
6. Thilagavathy, K.P, Santha, A. '*A Note on Achromatic and b-chromatic number of graphs*', International Journal of Applied Mathematics and Statistics, 53(1),104-110, 2016.
7. Thilagavathy, K.P, Santha, A. '*An Achromatic Number and b-chromatic number of Some Star related graphs*' Proceedings of National conference on Developing Scenario in Applied Sciences and Communicative English, Kumaraguru College of Technology, 305-307, 2016.
8. Thilagavathy, K. P, Santha, A. '*The Achromatic and b-Chromatic Colouring of Central Graph of Book Graph and Shadow graph of Path graph*' International Journal of pure and applied mathematics, 113,1-9,2017.
9. Vernold Vivin, J., Thilagavathi, K. '*On Harmonious colouring of Central Graphs*', Far East. Math. Sci (FJMS), 2,189-197, 2006.