

Computation of Even-Odd Harmonious Labeling of Certain Family of Acyclic Graphs

M. Kalaimathi, B. J. Balamurugan

Abstract: Let $G(V, E)$ be a graph with p number of vertices and q number of edges. An injective function $f : V \rightarrow \{1, 3, 5, \dots, 2p - 1\}$ is called an even-odd harmonious labeling of the graph G if there exists an induced edge function $f^* : E \rightarrow \{0, 2, \dots, 2(q - 1)\}$ such that

- i) f^* is bijective function
- ii) $f^*(e = uv) = (f(u) + f(v)) \pmod{2q}$

The graph obtained from this labeling is called even-odd harmonious graph.

Keywords : Graphs, Even-Odd Harmonious Labeling, Injective Function, Bijective Function.

I. INTRODUCTION

Labeling of a graph is an interesting and potential area of research in graph theory. It deals how the vertices and edges of the graph are labeled through well defined Mathematical functions [1]. The recent advancements and applications of various labelings of graphs have been updated by Gallian [5]. In the year 1967, Rosa in [9] introduced the graph labeling concepts. N. Lakshmi Prasana et al., in the paper [7] listed out the applications of graph labeling. We refer the text books written by Harary [4] and D. B. West [11] for the concepts and terminologies in graph theory.

The harmonious graph were introduced [3] in the year 1980. Further Z. Liang et al., [8] introduced the odd harmonious graphs in the year 2009 and P.B. Sarasija et al., [10] introduced the even harmonious graphs in the year 2011. Subsequently in the year 2015, Adalin Beatress and Sarasija in [2] introduced the even-odd harmonious graphs. Following this in the year 2019, we have proved that the graphs which are obtained through certain graph operations admit the even-odd harmonious labeling in the paper [6]. In this paper, we prove further the existence of even-odd harmonious labeling to certain family of acyclic graphs.

II. PRELIMINARIES

In this section, we recall the definitions of certain graphs pertaining to this paper.

Revised Manuscript Received on July 22, 2019.

* Correspondence Author

M. Kalaimathi *, Mathematics Division, School of Advanced Sciences, VIT University, Chennai, India. Email: kalaimathi.m2018@vitstudent.ac.in

B. J. Balamurugan, Mathematics Division, School of Advanced Sciences, VIT University, Chennai, India. Email: balamurugan.bj@vit.ac.in

Definition 2.1

A caterpillar is a graph obtained from a path graph by adding one or more pendant edges to the vertices of the path. The caterpillars can be drawn as a bipartite graph in a zigzag vertical fashion with one partitioned set on the left side and the other partitioned set on the right side. A caterpillar with m vertices on the path and t pendant edges is denoted $Cat_m^{+t}(l, r)$, where the vertex set of the caterpillar can be partitioned into two sets with l and r vertices.

Definition 2.2

1-regular lobster is a graph obtained from a path of n vertices P_n by attaching a path of length 2 to each vertex of P_n .

Definition 2.3

Let P_n be a path with n vertices. A coconut tree $CT_{n,m}$ is a graph obtained by joining m new pendent edges at an end vertex of P_n .

Definition 2.4

A tree T with n legs and l length is called spider tree if exactly one vertex of degree greater than or equal to 3.

Definition 2.5

A complete bipartite graph $K_{1,n}$ is called a star graph S_n and it has $n + 1$ vertices and n edges.

III. EVEN-ODD HARMONIOUS LABELING OF GRAPHS

In this section, we provide the notion of even-odd harmonious labeling of graphs [2].

Definition 3.1

Let $G(V, E)$ be a graph with p number of vertices and q number of edges. An injective function $f : V \rightarrow \{1, 3, 5, \dots, 2p - 1\}$ is called an even-odd harmonious labeling of the graph G if there exists an induced edge function $f^* : E \rightarrow \{0, 2, \dots, 2(q - 1)\}$ such that

- i) f^* is bijective function
- ii) $f^*(e = uv) = (f(u) + f(v)) \pmod{2q}$

Definition 3.2

A graph obtained by assigning numbers to the vertices and edges through an even-odd harmonious labeling is known as even-odd harmonious graph.

Remark

We use the phrase “EOH labeling of graph” instead of the phrase “even-odd harmonious labeling of graph” in this paper for simplicity.

Example 3.1




Fig. 1 EOH labeling of G

IV. EVEN-ODD HARMONIOUS LABELING OF ACYCLIC GRAPHS

In this section, we prove certain family of acyclic graphs admit the EOH labeling.

Theorem 4.1

The caterpillar graph $cat_m^{+t}(l,r)$ admits an EOH labeling when $m \geq 3, t \geq 1$.

Proof

Let $V = \{l_i : 1 \leq i \leq l\} \cup \{r_j : 1 \leq j \leq r\}$ be the vertex set of $cat_m^{+t}(l,r)$ where the vertices of the left side are l_i and the vertices of right side are r_j . Let $E = \{e_{ij} = l_i r_j : 1 \leq i \leq l, 1 \leq j \leq r\}$ be the edge set of the caterpillar graph $cat_m^{+t}(l,r)$. Here the caterpillar graph has $p = m + t$ vertices and $q = m + t - 1$ edges.

Define an injective function $f : V \rightarrow \{1, 3, \dots, 2(m+t) - 1\}$ such that

$$f(l_i) = 2i - 1, 1 \leq i \leq l$$

$$f(r_j) = 2l + 2j - 1, 1 \leq j \leq r$$

and an induced edge function $f^* : E \rightarrow \{0, 2, 4, \dots, 2(m+t-1) - 2\}$ such that

$$f^*(e_{ij}) = f^*(l_i r_j) = (2l + 2i) \pmod{2q},$$

$$1 \leq i \leq l, 1 \leq j \leq r$$

where f^* is bijective. The functions f and f^* give EOH labeling of G .

Therefore, the caterpillar $cat_m^{+t}(l,r)$ is an EOH graph when $n \geq 1$.

Example 4.1

An EOH labeling of $cat_8^{+5}(6,7)$ is shown in Fig. 2




Fig. 2 EOH labeling of $cat_8^{+5}(6,7)$

Theorem 4.2

The 1-regular lobster graph admits an EOH labeling when $n \geq 2$.

Proof

Let $V = \{u_i : 1 \leq i \leq n\} \cup \{u_{ij} : 1 \leq i \leq n \text{ and } j = 1, 2\}$ be the vertex set of 1-regular lobster where u_i are the vertices of the path P_n and u_{ij} are the vertices, which are adjacent to u_i for $1 \leq i \leq n$ and $j = 1, 2$.

Let $E = \{e_i = u_i u_{i+1} : 1 \leq i \leq n - 1\} \cup \{e_{ij} = u_i u_{ij} : 1 \leq i \leq n \text{ and } j = 1, 2\}$ be the edge set of the 1-regular lobster. Here the 1-regular lobster has $p = 3n$ vertices and $q = 3n - 1$ edges.

Define an injective function $f : V \rightarrow \{1, 3, \dots, 2(3n) - 1\}$ such that

Example 4.2

An EOH labeling of 1-regular lobster is shown in Fig. 3




Fig. 3 EOH labeling of 1-regular lobster

Case (i) $n \equiv 0 \pmod{2}$

$$f(u_i) = \begin{cases} 3(i-1)+1, & \text{if } i \text{ is odd} \\ 3n+4+3(i-2)+1, & \text{if } i \text{ is even} \end{cases}$$

$$f(u_{i1}) = \begin{cases} 3(i-1)+3n+1, & \text{if } i \text{ is odd} \\ 3(i-2)+5, & \text{if } i \text{ is even} \end{cases}$$

$$f(u_{i2}) = \begin{cases} 3(i-1)+3, & \text{if } i \text{ is odd} \\ 3(n+i)-3, & \text{if } i \text{ is even} \end{cases}$$

Case (ii) $n \equiv 1 \pmod{2}$

$$f(u_i) = \begin{cases} 3(i-1)+1, & \text{if } i \text{ is odd} \\ 3(n+i), & \text{if } i \text{ is even} \end{cases}$$

$$f(u_{i1}) = \begin{cases} 3(n+i)-1, & \text{if } i \text{ is odd} \\ 3(i-2)+5, & \text{if } i \text{ is even} \end{cases}$$

$$f(u_{i2}) = \begin{cases} 3(i-1)+3, & \text{if } i \text{ is odd} \\ 3(n+i-1)+1, & \text{if } i \text{ is even} \end{cases}$$

and an induced edge function

$$f^* : E \rightarrow \{0, 2, 4, \dots, 2(3n-1)-2\}$$
 such that

Case (i) $n \equiv 0 \pmod{2}$

$$f^*(e_i) = f^*(u_i u_{i+1}) = (3n+4+6(i-1)+2) \pmod{2q}, \quad 1 \leq i \leq n-1$$

$$f^*(e_{i1}) = f^*(u_i u_{i1}) = \begin{cases} (3n+6(i-1)+2) \pmod{2q}, & \text{if } i \text{ is odd} \\ (3n+6i-2) \pmod{2q}, & \text{if } i \text{ is even} \end{cases}$$

$$f^*(e_{i2}) = f^*(u_i u_{i2}) = \begin{cases} (3n+6(i-1)+4) \pmod{2q}, & \text{if } i \text{ is odd} \\ (3(n+2i-2)+2) \pmod{2q}, & \text{if } i \text{ is even} \end{cases}$$

Case (ii) $n \equiv 1 \pmod{2}$

$$f^*(e_i) = f^*(u_i u_{i+1}) = (3(n+2i)+1) \pmod{2q}, \quad 1 \leq i \leq n-1$$

$$f^*(e_{i1}) = f^*(u_i u_{i1}) = \begin{cases} (3(n+2i-1)-1) \pmod{2q}, & \text{if } i \text{ is odd} \\ (3(n+2i+1)+2) \pmod{2q}, & \text{if } i \text{ is even} \end{cases}$$

$$f^*(e_{i2}) = f^*(u_i u_{i2}) = \begin{cases} (3n+6(i-1)+5) \pmod{2q}, & \text{if } i \text{ is odd} \\ (3(n+2i-2)+3) \pmod{2q}, & \text{if } i \text{ is even} \end{cases}$$

where f^* is bijective. Here in all cases f and f^* defines the EOH labeling of G . Thus, the 1-regular lobster admits EOH labeling when $n \geq 2$.

Theorem 4.3

The coconut tree graph $CT_{n,m}$ admits an EOH labeling when $n \geq 2, m \geq 2$.

Proof

Let $V = \{u_i : 1 \leq i \leq n\} \cup \{v_j : 1 \leq j \leq m\}$ be the vertex set of coconut tree where u_i are the vertices of the path P_n and v_j are the m new pendent vertices at an end vertex of the path P_n .

Let $E = \{e_i = u_i u_{i+1} : 1 \leq i \leq n\} \cup \{e_{ij} = u_i v_j : i = n, 1 \leq j \leq m\}$ be the edge set of coconut tree. Here the coconut tree has $p = n + m$ vertices and $q = n + m - 1$ edges.

Define an injective function $f : V \rightarrow \{1, 3, \dots, 2(n+m)-1\}$ such that

Case (i) $n \equiv 1 \pmod{2}$

$$f(u_i) = \begin{cases} i, & \text{if } i \text{ is odd} \\ n+i, & \text{if } i \text{ is even} \end{cases}$$

$$f(v_j) = 2n+2j-1, \quad 1 \leq j \leq m$$

Case (ii) $n \equiv 0 \pmod{2}$

$$f(u_{2i-1}) = n+i, \quad 1 \leq i \leq \frac{n}{2}$$

$$f(u_{2i}) = 2i-1, \quad 1 \leq i \leq \frac{n}{2}$$

$$f(v_j) = 2n+2j-1, \quad 1 \leq j \leq m$$

and an induced edge function

$$f^* : E \rightarrow \{0, 2, 4, \dots, 2(n+m-1)-2\}$$
 such that

Case (i) $n \equiv 1 \pmod{2}$

$$f^*(e_i) = f^*(u_i u_{i+1}) = (n+1+2i) \pmod{2q}, 1 \leq i \leq n-1$$

$$f^*(e_{ij}) = f^*(u_i v_j) = (2(m+n-1)+2j) \pmod{2q}, i=n, 1 \leq j \leq m$$

Case (ii) $n \equiv 0 \pmod{2}$

$$f^*(e_i) = f^*(u_i u_{i+1}) = (n+2i) \pmod{2q}, 1 \leq i \leq n$$

$$f^*(e_{ij}) = f^*(u_i v_j) = (2n+2j+2) \pmod{2q}, i=n, 1 \leq j \leq m$$

where, f^* is bijective. The functions f and f^* provides the numbers on vertices and edges satisfying the conditions of EOH labeling of G . Thus, the coconut tree $CT_{n,m}$ admits EOH labeling when $n \geq 2, m \geq 2$.

Example 4.3

An EOH labeling of $CT_{3,5}$ and $CT_{4,5}$ is shown in Fig. 4 and Fig. 5




Fig. 4 EOH labeling of $CT_{3,5}$




Fig. 5 EOH labeling of $CT_{4,5}$

Theorem 4.4

The spider tree graph with n legs and l length admits an EOH labeling when $n \geq 3, n \equiv 1 \pmod{2}$ and $l \equiv 0 \pmod{2}$.

Proof

Let $V = \{v\} \cup \{v_{ij} : 1 \leq i \leq n, 1 \leq j \leq l\}$ be the vertex set of spider tree where v is a center vertex, v_{ij} are the n legs and l length of the spider tree graph. Let $E = \{e_{ij} = v_{ij} v_{ij+1} : 1 \leq i \leq n \text{ and } 1 \leq j \leq l-1\} \cup \{e_i = v v_{i1} : 1 \leq i \leq n\}$ be the edge set of the spider tree. Here the spider tree has $p = nl + 1$ vertices and $q = nl$ edges.

Define an injective function $f : V \rightarrow \{1, 3, \dots, 2(nl+1)-1\}$ such that

$$f(v) = 1$$

$$f(v_{ij}) = \begin{cases} l(n+i-1) + j + 2, & \text{if both } i \text{ and } j \text{ are odd} \\ l(n+i-1) + j + 1, & \text{if both } i \text{ and } j \text{ are even} \\ (i-1)l + j + 1, & \text{if } i \text{ is odd and } j \text{ is even} \\ (i-1)l + j + 2, & \text{if } i \text{ is even and } j \text{ is odd} \end{cases}$$

and an induced edge function $f^* : E \rightarrow \{0, 2, 4, \dots, 2nl-2\}$ such that

$$f^*(e_{ij}) = f^*(v_{ij} v_{ij+1}) = (l(n+3i-2) + 2(j-1)) \pmod{2q}, 1 \leq i \leq n \text{ and } 1 \leq j \leq l-1$$

$$f^*(e_i) = f^*(v v_{i1}) = \begin{cases} (l(n+i-1) + 3) \pmod{2q}, & \text{if } i \text{ is odd} \\ (l(i-1) + 3) \pmod{2q}, & \text{if } i \text{ is even} \end{cases}$$

where f^* is bijective. Both the functions f and f^* provides numbers to the vertices and edges satisfying EOH labeling of G . Therefore, the spider tree with n legs and l length admits EOH labeling.



Example 4.4

An EOH labeling of the spider tree with 5 legs and length 4 is shown in Fig. 6




Fig. 6 EOH labeling of spider tree with 5 legs and length 4

Theorem 4.5

The star graph $S_{m,3}$ admits an EOH labeling when $m \equiv 1 \pmod{2}$.

Proof

Let $V = \{x\} \cup \{u_i : 1 \leq i \leq m\} \cup \{v_i : 1 \leq i \leq m\} \cup \{w_i : 1 \leq i \leq m\}$ be the vertex set of star graph where x is a center vertex, u_i, v_i, w_i are the vertices of the path P_3 for $1 \leq i \leq m$. Let $E = \{e_i = uu_i : 1 \leq i \leq m\} \cup \{g_i = u_i v_i : 1 \leq i \leq m\} \cup \{h_i = v_i w_i : 1 \leq i \leq m\}$ be the edge set of the star graph $S_{m,3}$. Here the star graph has $p = 3m + 1$ vertices and $q = 3m$ edges.

Define an injective function $f : V \rightarrow \{1, 3, \dots, 2(3m + 1) - 1\}$ such that

$$f(x) = 2m + 1$$

$$f(u_1) = 6m + 1$$

$$f(u_{2i}) = 4i - 1, 1 \leq i \leq \frac{m}{2}$$

$$f(u_{2i+1}) = 2m + 4i + 1, 1 \leq i \leq \frac{m}{2}$$

$$f(v_{2i}) = 2m + 4i - 1, 1 \leq i \leq \frac{m}{2}$$

$$f(v_{2i-1}) = 4(i - 1) + 1, 1 \leq i \leq \frac{m}{2}$$

$$f(w_1) = 4m + 1$$

$$f(w_{2i}) = 5m + 2 - 2i, 1 \leq i \leq \frac{m}{2}$$

$$f(w_{2i+1}) = 6m - 2i + 1, 1 \leq i \leq \frac{m}{2}$$

and an induced edge function

$f^* : E \rightarrow \{0, 2, 4, \dots, 2(3m) - 2\}$ such that

$$f^*(e_1) = f^*(uu_1) = (2(m + 1)) \pmod{2q}$$

$$f^*(e_{2i}) = f^*(uu_{2i}) = (2m + 4i) \pmod{2q}, 1 \leq i \leq \frac{m}{2}$$

$$f^*(e_{2i+1}) = f^*(uu_{2i+1}) = (4m + 4i + 2) \pmod{2q}, 1 \leq i \leq \frac{m}{2}$$

$$f^*(g_1) = f^*(u_1 v_1) = 2$$

$$f^*(g_{i+1}) = f^*(u_{i+1} v_{i+1}) = (2(m + 1) + 4i) \pmod{2q}, 1 \leq i \leq m$$

$$f^*(h_1) = f^*(v_1 w_1) = (4m + 2) \pmod{2q}$$

$$f^*(h_{2i}) = f^*(v_{2i} w_{2i}) = (m + 2i + 1) \pmod{2q}, 1 \leq i \leq \frac{m}{2}$$

$$f^*(h_{2i+1}) = f^*(v_{2i+1} w_{2i+1}) = (2 + 2i) \pmod{2q}, 1 \leq i \leq \frac{m}{2}$$

where f^* is bijective. The functions f and f^* give EOH labeling of G . Thus, the star $S_{m,3}$ admits EOH labeling when $m \equiv 1 \pmod{2}$.

Example 4.5

An EOH labeling of $S_{5,3}$ is shown in Fig. 7




Fig. 7 EOH labeling of $S_{5,3}$

V. CONCLUSION

In this paper, we have proved that the family of acyclic graphs such as caterpillar, 1-regular lobster graph, coconut tree, spider tree and star graph admit the EOH labeling.

FUTURE SCOPE

We can find a family of cyclic graphs which will admit this EOH labeling.

REFERENCES

1. B.D. Acharya , S.M Hegde, Arithmetic Graphs. J. Graph Theory, Vol. 14, No.3, 275-299, (1990).
2. N. Adalin Beatress, P.B Sarasija, Even-Odd Harmonious Graphs, International Journal of Mathematics and Soft Computing, Vol. 5, No. 1, 23-29, (2015).
3. R.L Graham, N.J.A Sloane, On Additive Bases and Harmonious Graphs, SIAM Journal on Algebraic and Discrete Methods, Vol. 1, No. 4, 382-404, (1980).
4. F. Harary, Graph Theory, Addison-Wesley, Reading Mass, (1972).
5. Joseph A. Gallian, A Dynamic Survey of Graph Labeling, The Electronic Journal of Combinatorics, DS6, (2018).
6. M.Kalaimathi, B.J Balamurugan, Computation of Even-Odd Harmonious Labeling of Graphs Obtained by Graph Operations, Recent Trends in Pure and Applied Mathematics, AIP Conf. Proc. 2177, 020030-1 - 020030-9, (2019).
7. N. Lakshmi Prasana, K. Saravanthi, Nagalla Sudhakar, Applications of Graph Labeling in Major Areas of Computing Science. International Journal of Research in Computer and Communication Technology, Vol. 3, No. 8, (2014).
8. Z. Liang, Z. Bai Z, On The Odd Harmonious Graphs with Applications, J. Appl. Math. Comput., 29, 105-116, (2009).
9. A. Rosa, On Certin Valuations of the Vertices of a Graph, In Theory of Graphs(Internat.Sympos. Rome. 1966), Gordan and Breach, Newyork, Dunod, Paris, 349-359, (1967).
10. P. B. Sarasija, R. Binthiya, Even Harmonious Graphs with Applications, International Journal of Computer Science and Information Security, Vol. 9, No. 7, 161-163, (2011).
11. D. B. West, Introduction to Graph Theory, PHI Learning Private Limited, 2nd Edition, (2009).

AUTHORS PROFILE



M.Kalaimathi, received her M.Sc. and M.Phil. degree in Mathematics from Madras Christian College, University of Madras, Chennai, India. She is currently pursuing Ph.D. in Mathematics at VIT University, Chennai Campus, Chennai, India. Her area of research is graph theory.



Dr. B.J.Balamurugan, received his Ph.D degree in Mathematics from the University of Madras, Chennai, India. Currently, he is an Assistant Professor(Senior) of Mathematics in the School of Advanced Sciences at VIT University, Chennai Campus, Chennai, India. He has more than 22 years of teaching experience at

Undergraduate and Postgraduate level courses. Dr.B.J.Balamurugan has published more than 32 research papers in various journals and conference proceedings. His research interest includes graph theory, graph grammars, fuzzy logic and Petri nets.