


# Polyherbal formulation for Kidney and Liver Protection

S J Kabilan, R Baskar, G Poorani



**Abstract:** The objectives of the study were to assess the potential of herbal formulation made up of *Wedeliachinensis* leaves and *Boerhaaviadiffusa* roots in possessing activities like nephroprotectivity and hepatoprotectivity. Nephroprotective and hepatoprotective activity was evaluated using MTT cytotoxicity assay using mammalian cell culture. The results of the hepatoprotective and nephroprotective activities showed that the formulation mixture of herbs *Wedeliachinensis* and *Boerhaaviadiffusa* roots is an excellent source of organ stimulator with high therapeutical importance. The hepatoprotective and nephroprotective properties make this formulation a unique one focusing on liver and kidney diseases.

**Keywords:** Hepatoprotectivity, Nephroprotectivity, Polyherbal formulation

## I. INTRODUCTION

Nephroprotectivity is an activity of any compound protecting the kidney cells and functions [2]. Likewise, hepatoprotectivity was to protect the liver cells from harmful substances [7]. The above mentioned herbal plants contain few phytoconstituents that possess the ability to act as a nephroprotectant and hepatoprotectant [10]. They play a major role in maintenance of these organs and preventing them from getting damaged [19]. They also boost up their performance and have a cleansing activity on them [20]. They prevent aging of cells present in those organs keeping them healthy for longer time [21].

*Wedeliachinensis* (Manjalkarisalai in Tamil), Asteraceae is a well reputed herbal medicine in Siddha, Ayurvedha and Unani system of traditional medicine. Recent studies show the presence of diterpenes, flavanoids, triterpenes, phytosteroids and saponins. It is also reported to possess anti-inflammatory, antioxidant, analgesic, hepatoprotective, antimicrobial, CNS depressant, wound healing, antistress and anticancer activity [5].

*Boerhaaviadiffusa* (Mookirattaikeerai in Tamil) is one of the well known medicinal plants that are used to treat variety of human diseased conditions as mentioned in Ayurveda, Charaka Samhita, and Sushrita Samhita. Huge variety of phytochemicals like flavonoids, alkaloids, glycosides, rotenoids, steroids, triterpenoids, lipids, lignans, carbohydrates, proteins, and glycoproteins etc have been

reported from the herb. The promising therapeutic effects of this plant include diuretic, hepatoprotective, anti-inflammatory, anti-cancer, anti-fibrinolytic, immuno-modulatory, anti-diabetic, immuno-suppressive, analgesic, anti-lymphoproliferative and used for the treatment of TB [11].

The aim of this study is to develop an herbal formulation that possesses nephroprotective and hepatoprotective activity.

## II. MATERIALS AND METHODS

### A. Chemicals

DMEM medium, Fetal Bovine Serum (FBS), Trypsin, Saline, H<sub>2</sub>O<sub>2</sub>.

### B. Sample Collection and Extraction

#### Sample Collection

*Wedeliachinensis* leaves and *Boerhaaviadiffusa* root powder were collected from an FSSAI approved herbal powder manufacturer from Coimbatore, Tamil Nadu. All herbs were stored in air-tight, light resistant container for further use. The samples were labeled as the *Wedeliachinensis* leaves (WC), *Boerhaaviadiffusa* (BD) and Formulation mix.

#### Sample extraction

About 20g of powdered mix of these herbs was successively extracted with 150 ml of distilled water. Then it is allowed to evaporate in open air to obtain aqueous extracts. The extracts were filtered using membrane filter.

### C. Determination of Hepatoprotective activity

Human liver HepG2 cells were exposed to a medium containing H<sub>2</sub>O<sub>2</sub>(1mM) along with /without various concentrations of the formulation (100, 200, 300, 400 and 500 mg/ml). Then cytotoxicity was assessed by estimating the viability of HepG2 cells by MTT reduction assay. HepG2 cells were grown in DMEM culture medium and made into single-cell suspension and seeded into a 96-well flat bottom plate with 1 × 10<sup>4</sup> cells per well. After 48 hr incubation, 100 μL of 1mM H<sub>2</sub>O<sub>2</sub> was added to each well followed by the addition of 100 μL diluted extract at varying concentrations to the appropriate wells and the plates were incubated for further 48 hr at 37°C in a humidified incubator with 5% CO<sub>2</sub>. Supernatant was removed from each well, and 100 μL of MTT (0.5 mg/mL) was added. MTT enters the cell's mitochondria, where it is reduced to an insoluble, colored (dark purple) formazan product. The cells were then solubilized with 100 μL of an organic solvent DMSO and the released, solubilized formazan reagent is measured spectrophotometrically.

Manuscript published on 30 December 2019.

\* Correspondence Author (s)

S J Kabilan\*, Department of Biotechnology, Kalasalingam Academy of Research and Education, Krishnankoil, Tamilnadu, India. Email: sjkabil@gmail.com

R Baskar, Department of Biotechnology, Kumaraguru College of Technology, Coimbatore, India. Email: baskar.r.bt@kct.ac.in

G Poorani, Department of Biotechnology, Kumaraguru College of Technology, Coimbatore, India. Email: gpoorani.biotech@gmail.com

© The Authors. Published by Blue Eyes Intelligence Engineering and Sciences Publication (BEIESP). This is an open access article under the CC-BY-NC-ND license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

## Polyherbal formulation for Kidney and Liver Protection

Hepatoprotective activities of the formulation extracts are measured against the toxicity caused by H<sub>2</sub>O<sub>2</sub> on the liver cells and the readings were obtained at 540 nm (Surendran et al., 2011).

### D. Determination of in-vitro nephroprotective activity

Fresh African green monkey normal kidney cells (Vero) were grown in DMEM culture medium and made into single-cell suspension and seeded into a 96-well flat bottom plate with 1 × 10<sup>4</sup> cells per well. After 24 hr incubation, 100 μL of 1mM H<sub>2</sub>O<sub>2</sub> was added to each well followed by the addition of 100 μL diluted extract at varying concentrations to the appropriate wells and the plates were incubated for further 24 hr at 37°C in a humidified incubator with 5% CO<sub>2</sub>. Supernatant was removed from each well, and 100 μL of MTT (0.5 mg/mL) was added. MTT enters the cell's mitochondria, where it is reduced to an insoluble, colored (dark purple) formazan product. The cells are then solubilized with 100 μL of an organic solvent DMSO and the released, solubilized formazan reagent is measured spectrophotometrically. Nephroprotective activities of the formulation extracts are measured against the toxicity caused by H<sub>2</sub>O<sub>2</sub> on the liver cells and the readings were obtained at 540 nm (Srinivasan et al., 2015).

## III. RESULTS AND DISCUSSION

### A. Determination of Hepatoprotective activity

Hepatotoxicity was induced by H<sub>2</sub>O<sub>2</sub> in HepG2 cell lines. Formulation extract has been evaluated for its hepatoprotectiveness. The presence of phytoconstituents in formulation mix extract has inhibited the induced hepatotoxicity.

Table I shows the % of cell death occurred due to toxicity induced by H<sub>2</sub>O<sub>2</sub> with or without different concentration of Formulation extracts.

**Table I: Percentage cell death upon H<sub>2</sub>O<sub>2</sub> toxicity in HepG2 cell lines**

S.No	Test Concentration (μg/ml)	% of cell death in presence of formulation + H <sub>2</sub> O <sub>2</sub>
1	100	66.27 ± 2.68
2	200	54.25 ± 2.63
3	300	49.85 ± 3.51
4	400	43.98 ± 5.3
5	500	35.19 ± 5.85
H <sub>2</sub> O <sub>2</sub> toxic control		88.56 ± 1.75


The above data shows that with increasing concentration of formulation, the cell death has been reduced, which shows the protective activity of the formulation.

Table II: Hepatoprotective activity of formulation mix over toxicant in HepG2 cell line

S.No	Test Concentration (μg/ml)	% Protection offered over toxicant control
1	100	25.16 ± 3.03
2	200	38.74 ± 2.98
3	300	43.70 ± 3.97
4	400	50.33 ± 5.98

5	500	60.26 ± 6.61
---	-----	--------------

Figure 1: Percentage hepatoprotectivity of formulation mix over toxic control



The PC<sub>50</sub> (Protective Concentration 50%) is 381.48 μg/ml which results in better hepatoprotective activity over the H<sub>2</sub>O<sub>2</sub> toxicity in HepG2 cells.

It has been revealed from Table II, that effective protection of about 50 % over toxicant control has been attained at 381.48 μg/ml concentration (Srinivasan et al., 2015). Both the *Boerhaaviadiffusa* and *Wedeliachinensis* possess hepatoprotective activity and contains phytoconstituents having hepatoprotective activity.

### B. Determination of nephroprotective activity

Nephrotoxicity was induced by H<sub>2</sub>O<sub>2</sub> in Vero cell lines. Formulation mix extract has been evaluated for its nephroprotectiveness. The presence of various phytoconstituents and minerals in formulation mix extract has inhibited the induced nephrototoxicity (Ahmed et al., 2010).

Table III shows the % of cell death occurred due to toxicity induced by H<sub>2</sub>O<sub>2</sub> with or without different concentration of formulation mix extracts.

Table III: Percentage cell death upon H<sub>2</sub>O<sub>2</sub> toxicity in Vero cell lines


S.No	Test Concentration (μg/ml)	% of cell death in presence of formulation + H <sub>2</sub> O <sub>2</sub>
1	100	89.63 ± 0.44
2	200	88.34 ± 0.77
3	300	83.41 ± 0.44
4	400	67.09 ± 1.18
5	500	63.21 ± 1.18
H <sub>2</sub> O <sub>2</sub> toxic control		93.26 ± 0.44

From the table, it is evident that the H<sub>2</sub>O<sub>2</sub> toxicity was inhibited with increasing concentration of formulation.

Table IV: Nephroprotective activity of formulation mix over toxicant in Vero cell line

S.No	Test Concentration (µg/ml)	% Protection offered over toxicant control
1	100	3.89 ± 0.48
2	200	5.28 ± 0.83
3	300	10.56 ± 0.48
4	400	28.06 ± 1.27
5	500	32.22 ± 1.28

Figure 2: % nephroprotectivity of formulation mix over toxic control



The PC50 (Protective Concentration 50%) is 732.06µg/ml

It has been shown from Table IV, that effective protection of about 50 % over toxicant control has been attained at 732.06µg/ml concentration (Kiruba et al., 2014). *Boerhaaviadiffusa* possess nephroprotective activity and contains phytoconstituents having nephroprotective activity which results in moderate nephroprotective activity over the H<sub>2</sub>O<sub>2</sub> toxicity in Vero cells.

#### IV. CONCLUSION

The results of hepatoprotective and nephroprotective activities showed that the formulation mixture of herbs *Wedeliachinensis* and *Boerhaaviadiffusa* roots is an excellent source of organ protector with high therapeutical importance. The hepatoprotective and nephroprotective properties make this formulation a unique one focusing on liver and kidney diseases. It may therefore be recommended for people to prevent and get rid of kidney and liver related disorders.

#### ACKNOWLEDGMENT

The authors acknowledge Department of Biotechnology and Re forum (Research Cell) of KCT for providing essential facilities and funding to perform the research.

#### REFERENCES

[1] Ahmed, Mahgoub M. (2010). "Biochemical studies on nephroprotective effect of carob (*Ceratonia siliqua* L.) growing in Egypt." *Nature and Science*, Vol. 8, 3: 41-47.

[2] Alshawsh, M. A., Abdulla, M. A., Ismail, S., & Amin, Z. A. (2011). Hepatoprotective effects of *Orthosiphon stamineus* extract on thioacetamide-induced liver cirrhosis in rats. *Evidence-Based Complementary and Alternative Medicine*, 2011.

[3] Fotakis, G., & Timbrell, J. A. (2006). In vitro cytotoxicity assays: comparison of LDH, neutral red, MTT and protein assay in hepatoma cell lines following exposure to cadmium chloride. *Toxicology letters*, 160(2), 171-177.

[4] Hassan, S. W., Salawu, K., Ladan, M. J., Hassan, L. G., Umar, R. A., & Fatihu, M. Y. (2010). Hepatoprotective, antioxidant and phytochemical properties of leaf extracts of *Newbouldia laevis*. *International Journal of Pharm Tech Research*, 2(1), 573-584.

[5] Herberg, J., Pahari, A., Walters, S., & Levin, M. (2009). Infectious diseases and the kidney. In *Pediatric Nephrology*(pp. 1235-1273). Springer Berlin Heidelberg.

[6] Kanniparambil Xavier, S., Haneefa, S. M., & Anand, D. R. (2017). Antioxidant and nephroprotective activities of the extract and fractions of *Homonoia riparia* Lour. *Pharmacognosy magazine*, 13(49), 25.

[7] Kiruba, Arun K. P., Brindha P. (2014). In vitro studies on nephroprotective efficacy of *Cynodon dactylon* and *Gmelina asiatica*. *Asian journal of pharmaceutical and clinical research*, 7, 111-120.

[8] Koul, S., Pandurangan, A., & Khosa, R. L. (2012). *Wedelia chinensis* (Asteraceae)—an overview. *Asian Pacific Journal of Tropical Biomedicine*, 2(2), S1169-S1175.

[9] Lakshmi, M. S., Reddy, U. K., & Rani, S. R. K. S. (2012). A review on medicinal plants for nephroprotective activity. *Asian J Pharm Clin Res*, 5(4), 8-14.

[10] Mahesh, A. R., Kumar, H., Ranganath, M. K., & Devkar, R. A. (2012). Detail study on *Boerhavia Diffusa* plant for its medicinal importance-a review. *Res J Pharm Sci*, 1, 28-36.

[11] Mandal, S., & Bapna, J. S. (2016). Pharmacological evaluation of *Boerhavia diffusa* plant extract for their potential in management of diabetic nephropathy in Streptozotocin induced Type I diabetic rats 5(12), 1097-1108.

[12] Murti, K., Panchal, M. A., & Lambole, V. (2010). Pharmacological properties of *Boerhaavia diffusa* - a review. *Int J Pharm Sci Rev Res*, 5(2), 107-110.

[13] Nayak, P., & Thirunavoukkarasu, M. (2016). A review of the plant *Boerhaavia diffusa*: Its chemistry, pharmacology and therapeutical potential. *J. Phytopharmacol*, 5(2), 83-92.

[14] Nomani, I., Mazumder, A., & Chakraborty, G. S. (2013). *Wedelia chinensis* (Asteraceae)-An Overview Of A Potent Medicinal Herb. *Int J PharmTech Res*, 5(3), 957-964.

[15] Ramesh, K., Manohar, S., & Rajeshkumar, S. (2014). Nephroprotective activity of ethanolic extract of *Orthosiphon stamineus* leaves on ethylene glycol induced urolithiasis in albino rats. *J. PharmTech Res*, 6(1), 403-408.

[16] Reshma, A. K., & Brindha, P. (2014). In Vitro Anti-Inflammatory, Antioxidant and Nephroprotective Studies on Leaves of *Aegle Marmelos* and *Ocimum Sanctum*. *Asian journal of pharmaceutical and clinical research*, 7, 121-124.

[17] Sasidharan, S., Aravindran, S., Latha, L. Y., Vijenth, R., Saravanan, D., & Amutha, S. (2010). In vitro antioxidant activity and hepatoprotective effects of *Lentinula edodes* against paracetamol-induced hepatotoxicity. *Molecules*, 15(6), 4478-4489.

[18] Srinivasan, V., Panneerselvam, R., & Gunasekaran, S. (2015). Nephro-protective Activity of Ethanolic Extract of *Melia Azadirachta* against H<sub>2</sub>O<sub>2</sub>Induced Toxicity in Vero Cell Line.

[19] Sundararajan, R., Bharampuram, A., & Koduru, R. (2014). A review on phytoconstituents for nephroprotective activity. *Pharmacophore*, 5(1).

[20] Surendran, S., Eswaran, M. B., Vijayakumar, M., & Rao, C. V. (2011). In vitro and in vivo hepatoprotective activity of *Cissampelos pareira* against carbon-tetrachloride induced hepatic damage. Talwani, R., Gilliam, B. L., & Howell, C. (2011). Infectious diseases and the liver. *Clinics in liver disease*, 15(1), 111-130.

## AUTHORS PROFILE



**S J Kabilan**, completed his B.Tech (Biotechnology) at Kalasalingam University and M.Tech (Biotechnology) at Kumaraguru College of Technology (Affiliated to Anna University). Both the Degrees completed with First class with Distinction. Also, pursuing PhD in the area of Herbal Drug Research. Currently Employment: Assistant Professor, Department of Biotechnology, School of Bio and Chemical Engineering, Kalasalingam Academy of Research and Education, Tamilnadu, India.

**Previous Publications:** Kabilan, M. S. (2018). Antioxidant and anti-inflammatory properties of G-immune plus: A polyherbal formulation. *International Journal of Green Pharmacy (IJGP)*, 12(03).



**Dr. R. Baskar**, PhD in Medical Biochemistry. Currently working as Associate professor at Kumaraguru college of Technology, Coimbatore, Tamilnadu, India.

### Memberships:


Life Member in Indian Society for Technical Education  
Life Member in National Society for Ethnopharmacology.

### Achievements:

1. INSA Summer Fellowship award during 2012 to work in the Laboratory of Plant-Microbe Interaction, School of Botany, University of Hyderabad on "Plant Growth Promoting Rhizobacteria".
2. Reviewer in National and International Journals
3. Currently Academic Editor in Biotechnology Journal International.
4. No. of citations : 792; h index : 12; i10 index : 16

### Recent Publications:

- **Baskar, R.**, Akshaya, S.B., Akshitha, R., Dhilip Kumar, G., Poorani, G. (2018). Evaluation of antioxidant and phytochemical activity in solvent extracts of *Delonix regia*. *International Journal of Green Pharmacy*, Volume 12(3), July-September issue, S607-S616.
- **Baskar, R.**, Sweatha, S., Karunambika, R., G.Ramya, R., Shanthi (2017). Influence of methionine on ergothioneine content and antioxidant activity of *Pleurotus platypus*. *Asian Journal of Microbiology, Biochemistry and Environmental Sciences*, Volume 19 (Supplementary issue), S34-S38.
- Gurumallesh, P., Alagu, K., **Ramakrishnan, B.** and Muthusamy, S (2019). A systematic reconsideration on proteases. *International Journal of Biological Macromolecules*, 128, 254-267 (**Impact Factor: 4.7**)
- Poorani Gurumallesh, **Baskar Ramakrishnan** and Bharathi Dhurai (2019). A novel metalloprotease from banana peel and its biochemical characterization. *International Journal of Biological Macromolecules*, 134, 527- 535 (**Impact factor: 4.7**)



**Poorani G** completed her B.Tech (Biotechnology) at Tamil Nadu Agricultural University, Coimbatore and M.Tech Pharmaceutical Technology at SASTRA University, Thanjavur and currently pursuing Ph.D – Kumaraguru College of Technology.

### Notable Publications:

INTERNATIONAL JOURNAL OF BIOLOGICAL MACROMOLECULES (2019) IF 4.78 A novel metalloprotease from banana peel and its biochemical characterization

MATERIAL SCIENCE AND ENGINEERING C (2019) IF 5.08 Biological synergy of greener gold nanoparticles by using *Coleus aromaticus* leaf extract

INTERNATIONAL JOURNAL OF BIOLOGICAL MACROMOLECULES (2019) IF 4.78 A systematic reconsideration on proteases

PROCESS BIOCHEMISTRY (2019) IF 2.88 Green synthesis of anisotropic silver nanoparticles from the aqueous leaf extract of *Dodonaea viscosa* with their Antibacterial and Anticancer activities

JOURNAL OF PHOTOCHEMISTRY AND PHOTOBIOLOGY B: BIOLOGY (2018) IF 4.06 Improved Conductivity and Antibacterial activity of poly (2- aminothiophenol) - silver nanocomposite against human pathogens

JOURNAL OF MICROENCAPSULATION (2016) IF 2.04 Formulation, characterization, in vitro and in vivo evaluation of castor oil based self-nano emulsifying levosulpiride delivery systems