

# On $\Delta$ Generalised Star Semi-Closed Sets in Topological Spaces



S. Rajakumar

**Abstract:** A modern form of sets labeled  $\delta g^*s$ -closed sets is imported in this material. A part of utilization of  $\delta g^*s$ -closed sets and its resources are explained. A modernistic space called  $\delta g^*s$ - $T^*_{3/4}$ -space is also popularized.

**Keywords:**  $\delta$ -closed sets,  $\delta$ -open sets,  $\delta g^*s$ -open sets,  $\delta g^*s$ -closed sets and  $\delta g^*s$ - $T^*_{3/4}$ -space.

## I INTRODUCTION

In 1968, Velicko [13] introduced the  $\delta$  closed sets. Some authors([1], [4], [8], [10], [11]) continuing their studies on semi-closed sets and continuous maps on topological spaces.  $g^*s$ -closed sets were made known by the topologist [14].  $\delta g^*s$ -closed sets and  $\delta g^*s$ - $T^*_{3/4}$ -space are popularized in this note by the author.

## II PRELIMINARIES

In every place of this text  $(N, \tau)$  or  $N$  perform topological spaces. Definitions of  $g$ ,  $sg$ ,  $gs$ ,  $ag$ ,  $\delta g$ ,  $ag$ -closed sets are collected from [5], [2], [7], [6], [3], [12], [9], [14].

## III. MAIN RESULTS

**Definition 3.1** A division  $M$  of  $(N, \tau)$  is termed as  $\delta g^*s$ -closed if  $cl_\delta(M) \subseteq V$  whenever  $M \subseteq V$ ,  $V$  is  $g^*s$ -open.

**Theorem 3.2** Each one  $\delta$ -closed  $\Rightarrow \delta g^*s$ -closed set.

**Proof.** Approve  $U \subseteq \delta$ -closed,  $V \subseteq g^*s$ -open consisting of  $U$ . As  $cl_\delta(U) = U$  for each  $U$  of  $N$ . Thus  $cl_\delta(U) \subseteq U \Rightarrow U$  is  $\delta g^*s$ -closed.

**Note 3.3,**  $\delta g^*s$ -closed  $\Rightarrow \delta$ -closed.

**Example 3.4** Let  $N = \{m, n, o, p\}$  with the topology  $\tau = \{N, \emptyset, \{m\}, \{n\}, \{m, n\}, \{m, n, o\}\}$ . Though the set  $\{o, p\}$  is  $\delta g^*s$ -closed it is not  $\delta$ -closed.

**Manuscript published on 30 December 2019.**

\* Correspondence Author (s)

S. Rajakumar\*, Assistant Professor, Department of Mathematics, Kalasalingam Academy of Research And Education, Krishnankoil-626 126, India, Email: srkumar277@gmail.com

© The Authors. Published by Blue Eyes Intelligence Engineering and Sciences Publication (BEIESP). This is an open access article under the CC-BY-NC-ND license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Proposition 3.5**  $\delta g^*s$  closed  $\Rightarrow g$ -closed set.

**Proof.** Authorize  $U = \delta g^*s$ -closed,  $V =$  open set consists of  $U$ .

Here  $U$  is  $\delta g^*s$  closed,  $cl_\delta(U) \subseteq U$  for each  $U$  of  $N$ . As  $cl(U) \subseteq cl_\delta(U) \subseteq U$ ,  $cl(U) \subseteq U$ , consequently  $U$  is  $g$ -closed.

**Example 3.6** Let  $N = \{s, t, u, v\}$ ,  $\tau = \{N, \emptyset, \{s\}, \{t\}, \{s, t\}, \{t, u, v\}\}$ . Then  $\{v\}$  is  $g$ -closed set and  $\neq \delta g^*s$ -closed in  $N$ . This example proves that  $g$ -closed  $\neq \delta g^*s$ -closed.

**Theorem 3.7** Every  $\delta g^*s$ -closed  $\Rightarrow gs$ -closed.

**Proof.** Presume  $U = \delta g^*s$ -closed.  $V =$  open set consists of  $U$ . As  $cl(U) \subseteq U$  for every  $U \subseteq N$ . As  $scl(U) \subseteq cl_\delta(U) \subseteq V$ ,  $scl(U) \subseteq V$ ,  $\Rightarrow U$  is  $gs$ -closed.

**Example 3.8** Let  $N = \{\alpha, \kappa, \theta, \psi\}$ ,  $\tau = \{N, \emptyset, \{\alpha\}, \{\kappa\}, \{\alpha, \kappa\}, \{\kappa, \theta, \psi\}\}$ . Though  $\{\theta\}$  is  $gs$ -closed  $\neq \delta g^*s$ -closed. This shows that  $gs$ -closed  $\neq \delta g^*s$ -closed.

**Theorem 3.9**  $\delta g^*s$ -closed  $\Rightarrow ag$ -closed set.

**Proof.** For each  $U$  of  $N$ , It is clear that  $acl(U) \subseteq cl_\delta(U)$ .

**Example 3.10** Let  $N = \{1, 2, 3, 4\}$ ,  $\tau = \{N, \emptyset, \{1\}, \{2\}, \{1, 2\}, \{2, 3, 4\}\}$ . Then  $\{1, 3\}$  is  $ag$ -closed set  $\delta g^*s$ -closed. However converse part fails in Theorem 3.9.

**Theorem 3.11**  $\delta g^*s$ -closed  $\Rightarrow \delta g$ -closed set.

**Proof.** Suppose  $U \subseteq \delta g^*s$ -closed,  $V$  the open set consists of  $U$ . As  $cl_\delta(A) \subseteq U$ , when  $A \subseteq U$ ,  $U$  is  $g^*s$ -open. There upon  $cl_\delta(U) \subseteq U$ ,  $U$  is open. Thus  $U$  is  $\delta g$ -closed.


**Example 3.12** Take  $N = \{\chi, \psi, \gamma, \delta\}$ ,  $\tau = \{N, \emptyset, \{\chi\}, \{\psi\}, \{\chi, \psi\}, \{\psi, \gamma, \delta\}\}$ . Then  $\{\chi, \delta\}$  is  $\delta g$ -closed set  $\neq \delta g^*s$ -closed in  $N$ . It proves that confer gets blunder in Theorem 3.11.

**Theorem 3.13**  $\delta g^*s$  -closed  $\Rightarrow \alpha g^\wedge$ -closed set.

**Proof.** For every  $D$  of  $(L, \tau)$ ,  $\alpha cl(D) \subseteq cl_\delta(D)$  is clear.

**Example 3.14** Let  $N = \{ \eta, \kappa, \lambda, \mu \}$ , with the topology  $\tau = \{ N, \phi, \{ \eta \}, \{ \kappa \}, \{ \eta, \kappa \}, \{ \kappa, \lambda, \mu \} \}$ . Then  $\{ \eta, \kappa, \lambda \}$  is  $\alpha g^\wedge$ -closed set  $\neq \delta g^*s$  -closed in  $N$ . Hence exchange is not perfect in Theorem 3.13.

**Note 3.15** Examples for relationships of  $\delta g^*s$  -closed with more noted sets.



**Remark 3.16** Self-reliant results with  $\delta g^*s$  -closed are given below.

**Illustration 3.17** Assume  $N = \{ \eta, \sigma, \rho \}$ ,  $\tau = \{ N, \phi, \{ \eta \}, \{ \sigma \}, \{ \eta, \sigma \} \}$ . Then  $\{ \eta, \sigma \}$  is  $\delta g^*s$  -closed but  $\neq g^*s$  -closed and  $\neq sg$ -closed

**Example 3.18** Take  $N = \{ \gamma, \rho, \sigma, s \}$  with the topology  $\tau = \{ N, \phi, \{ \gamma \}, \{ \gamma, \rho \}, \{ \sigma, s \}, \{ \gamma, \sigma, s \} \}$ . Then  $\{ \gamma, \rho \}$  is  $\delta g^*s$  closed but  $\neq g^\alpha$  -closed.

**Illustration 3.19** Assume  $N = \{ \theta, \sigma, \omega, \psi \}$  with the topology  $\tau = \{ N, \phi, \{ \theta \}, \{ \sigma \}, \{ \theta, \sigma \}, \{ \sigma, \omega, \psi \} \}$ . Then  $\{ \theta, \omega \}$  is  $g^*s, sg, g^\alpha$ -closed and  $\neq \delta g^*s$  -closed.

**Illustration 3.20** Consider  $N = \{ \gamma, \eta, \pi, \rho \}$  with the topology  $\tau = \{ N, \phi, \{ \gamma \}, \{ \gamma, \eta \}, \{ \pi, \rho \}, \{ \gamma, \pi, \rho \} \}$ . Then  $\{ \gamma \}$  is  $\alpha$ -closed and  $\neq \delta g^*$ -closed set.

#### IV. CHARACTERIZATIONS

**Theorem 4.1** The limited combination of  $\delta g^*s$  -closed sets  $\rightarrow \delta g^*s$  closed.

**Proof.** Endorse  $\{ X_i / i=1,2,\dots,n \}$  be a limited set of  $\delta g^*s$  -closed subsets  $N$ . Moreover for  $N_i, cl_\delta(X_i) \subseteq U_i, i \in \{ 1, 2, \dots, n \}$ . Hence  $N_i \subseteq \bigcup U_i = V$ . Here  $U$  of  $g^*s$ -open =  $g^*s$ -open in  $N$ .  $V$  is  $g^*s$ -open in  $N$ . In addition,  $\bigcup cl_\delta(N_i) = cl_\delta(\bigcup N_i) \subseteq V$ . Therefore  $\bigcup N_i$  is  $\delta g^*s$  -closed in  $N$ .

**Observation 4.2**  $\cap$  of  $\delta g^*s$  -closed sets in  $L \neq \delta g^*s$  -closed set. In example 3.14  $\{ \eta, \kappa, \lambda \} \cap \{ \eta, \kappa, \mu \} = \{ \eta, \kappa \}$  is  $\neq \delta g^*s$  -closed.

**Theorem 4.3** If  $E$  is a  $\delta g^*s$  -closed  $\subset$  of  $N$ , then  $cl_\delta(E) - E$  not  $\subseteq$  an  $g^*s$  -closed set.

**Proof.** Take  $U$  as  $\delta g^*s$  -closed and assume  $G$  be a  $g^*s$  closed set contained in  $cl_\delta(U) - U$ . Now  $G^c = g^*s$  open set of  $N$ , likewise  $U \subset G^c$ . As  $U = \delta g^*s$  -closed set of  $N$ , then  $cl_\delta(U) \subseteq G^c$ . Thus  $G \subseteq (cl_\delta(U))^c$ . Also  $G \subseteq cl_\delta(U) - U$ . Therefore  $G \subseteq (cl_\delta(U))^c \cap (cl_\delta(U)) = \phi$ . Thus  $G = \phi$ .

**Theorem 4.4** In  $N, U = g^*s$  open,  $\delta g^*s$  -closed  $\subset$  of  $N$  then is  $\delta$ -closed  $\subset$  of  $N$ .

**Proof.** As  $g^*s$  -open and  $\delta g^*s$  -closed  $= U, cl_\delta(U) \subseteq U$ . Hence  $U$  is  $\delta$ -closed.

**Theorem 4.5** In  $T_{3/4}$ -space each one  $\delta g^*s$  -closed set  $\Rightarrow \delta$ -closed.

**Proof.** Consider  $U$  be  $\delta g^*s$  -closed set of  $N$ , where  $N$  is  $T_{3/4}$ -Space. Here, each one  $\delta g^*s$  -closed set  $\rightarrow \delta g$  closed. Therefore  $N$  is  $T_{3/4}$ -Space and  $U$  is  $\delta$ -closed.

**Theorem 4.6** In  $N, U$  is  $\delta$ -closed and  $\delta g^*s$  -closed if and only if  $cl_\delta(U) - U = g^*s$  -closed.

**Proof.** Necessity. Assume  $U$  be a  $\delta$ -closed  $\subset$  of  $N$ . Then  $cl_\delta(U) = U$  and so  $cl_\delta(U) - U = \phi$ , a  $g^*s$  -closed. Adequate: As  $U$  is  $\delta g^*s$  -closed by Theorem 4.3,  $cl_\delta(A) - A \neq g^*s$ -closed. But  $cl_\delta(U) - U = \phi$ . That is  $cl_\delta(U) = U$ . Hence  $U$  is  $\delta$ -closed.

#### V. APPLICATIONS

**Definition 5.1**  $N$  is termed as  $\delta g^*s - T^*_{3/4}$  -space if each  $\delta g^*s$ -closed set  $\Rightarrow \delta$ -closed.

**Theorem 5.2** Every  $T_{3/4}$ -space is a  $\delta g^*s - T^*_{3/4}$ -space.

**Proof.** Here each  $\delta g^*s$ -closed  $\Rightarrow \delta g$ -closed, the argument is clear.

**Remark 5.3**  $\delta g^*s - T^*_{3/4}$  -space  $\neq T^*_{3/4}$  -space.

**Illustration 5.4** Consider  $N = \{ \delta, \pi, \theta \}$  and  $\tau = \{ N, \phi, \{ \delta \} \}$ .  $N$  is a  $\delta g^*s - T^*_{3/4}$  -space but not a  $T^*_{3/4}$  -space.

**Theorem 5.5** Every  $\delta g^*s - T^*_{3/4}$  -space is a  $T_{\alpha g^\wedge}$  -space.

**Proof.** Concede  $N$  be a  $\delta g^*s - T^*_{3/4}$  -space, then each singleton is  $g^*s$  -closed or  $\delta$ -open. As each singleton is  $g^*s$  -closed or  $\alpha$ -open,  $N$  is a  $T_{\alpha g^\wedge}$  -space

**Note 5.6**  $T_{\alpha\beta}$ -space  $\neq \delta g^*s$  -  $T^{*3/4}$ -space.

**Illustration 5.7** Consider  $N = \{ \chi, \pi, \xi \}$  and  $\tau = \{ N, \varphi, \{ \chi \}, \{ \xi \}, \{ \chi, \xi \}, \{ \pi, \xi \} \}$ .  $N$  is a  $T_{\alpha\beta}$ -space but  $\neq \delta g^*s$  -  $T^{*3/4}$ -space.

**Remark 5.8**  $\delta g^*s$  -  $T^{*3/4}$ -space and  $T_{1/2}$ -space are self-reliant to one another

**Illustration 5.9** Take  $N = \{ \chi, \psi, \zeta \}$  and  $\tau = \{ N, \varphi, \{ \chi \} \}$ .  $(N, \tau)$  is  $\delta g^*s$  -  $T^{*3/4}$ -space and  $\neq T_{1/2}$ -space.

**Illustration 5.10** Assume  $N = \{ \lambda, \zeta, \theta \}$  and  $\tau = \{ N, \varphi, \{ \lambda \}, \{ \theta \}, \{ \lambda, \theta \}, \{ \zeta, \theta \} \}$ .  $N$  is a  $T_{1/2}$ -space and  $\neq \delta g^*s$  -  $T^{*3/4}$ -space

### ACKNOWLEDGMENT

The biographer desire to give thanks to the conciliators for their impressive commentary and verdict which will guide to develop this publication

### REFERENCES

1. S.P. Arya, T. Nour, Characterization of S-normal spaces. *Indian J. Pure Appl. Math.* 21(8)(1990)717-719.
2. P.Bhattacharya and B.K. Lahiri, Semi generalized closed sets in Topology, *Indian J. Math.*, 29(1987), 375-382.
3. J. Dontchev and M. Ganster, On  $\delta$ -generalized closed sets and  $T_{3/4}$ -spaces, *Mem. Fac. Sci. Kochi Univ. Ser. A, Math.*, 17(1996), 15-31.0
4. N. Levine, Semi-open sets and Semi-continuity in topological spaces *Amer Math. Monthly*, 70(1963), 36-41.
5. N. Levine, Generalized closed sets in topology *Rend. Circ. Mat. Palermo*, 19(1970)89-96.
6. H.Maki, R. Devi and K. Balachandran, Generalized  $\alpha$ -closed sets in topology, *Bull. Fukuoka Univ. Ed part III*, 42(1993), 13-21.
7. H. Maki, R. Devi and K. Balachandran, Associated topologies of generalized  $\alpha$ -closed sets and  $\alpha$ -generalized closed sets, *Mem. Fac. Sci. Kochi Univ. Ser. A. Math.*, 15(1994), 57-63.
8. A.S.Mashhour, M.E Abd El-Monsef and S.N.El-Debb, On precontinuous and weak precontinuous mappings, *Proc. Math. and Phys. Soc. Egypt* 55(1982), 47-53.
9. M.E Abd El-Monsef, S.Rose Mary and M. Lellis Thivagar, On  $\alpha G$ -closed sets in topological spaces, *Assiut University Journal of Mathematics and Computr science*, Vol 36(1), P-P.43-51(2007).
10. O. Njastad, On some classes of nearly open sets, *Pacific J. Math.*, 15(1965), 961-970.
11. M. Stone, Application of theory of Boolean rings to general topology, *Trans. Amer. Math. Soc.*, 41(1937), 374-481.
12. M. K. R. S. Veerakumar,  $g^{\wedge}$ -closed sets in topological spaces, *Bull. Allah. Math. Soc.*, 18(2003), 99-112.
13. N.V.Velico, H-closed topological spaces, *Amer. Math. Soc. Transl.* 78(1968), 103-118.
14. M. K. R. S. Veerakumar,  $g^*$ -semi closed sets, *Acta Ciencia Indica. Acta Ciencia Indica*, Vol-29 M, No.1, 081(2003), 81-90.
15. Indica, Vol-29 M, No.1, 081(2003), 81-90.

### AUTHORS PROFILE



**Dr. S. Rajakumar** is currently a Assistant Professor in the Department of Mathematics in Kalasalingam Academy of Research And Education (Deemed to be University), Krishnankoil-626126. He obtained his Ph.D., degree in Manonmaniam Sundaranar University, Tirunelveli. His area of research interest is Topology and Bitopological spaces. He has 17+ years of teaching experience. He has a few publications in standard journals.